

~~98/42.~~

I 3.42

R54936



MEDICAL FACTS

AND

OBSERVATIONS.



VOL. VI.

MEDICAL FACTS

AND

OBSERVATIONS.

VOLUME THE SIXTH.

L O N D O N :

PRINTED FOR J. JOHNSON, N° 72, ST. PAUL'S CHURCH YARD.

M.DCC.XCV.

Digitized by the Internet Archive
in 2016

C O N T E N T S.

	Page
I. <i>OBSERVATIONS on the Use of Arsenic in the Intermittent Fevers of a tropical Climate; to which is prefixed an Account of the Weather, at Sierra Leone, during the Season in which such Fevers are most prevalent. By Thomas Masterman Winterbottom, M. D. Physician to the Settlement at Sierra Leone.</i> — —	1
II. <i>An Account of the good Effects of a Solution of Sal Ammoniac, in Vinegar, employed, as a topical Application, in Cases of lacerated Wounds. By Mr. Henry Yates Carter, Surgeon at Kettley, near Wellington, in Shropshire.</i> — —	66
III. <i>Case of a diseased Kidney. By the same.</i>	85
IV. <i>Case of a Gun-Shot Wound of the Head. By the same.</i> — — —	91
V. <i>An Account of some extraordinary Symptoms which were apparently connected</i> <i>with</i>	

with certain morbid Alterations about the Veins and Nerves. By Mr. John Pearfon, Surgeon of the Lock Hospital, and of the Public Dispensary. — 96

VI. *An Account of the Extraction of an extraneous Substance from the Rectum. By Mr. William Blair, Surgeon of the Lock Hospital; and of the General Dispensary in Newman Street, St. Mary-le-bone.* III

VII. *A Case of Aneurism of the Crural Artery. By Mr. Thompson Forster, Surgeon on the Staff of the Army, and Surgeon to Guy's Hospital.* — II4

VIII. *An Account of a Key Instrument of a new Construction; with Observations on the Principles on which it acts, in the Extraction of Teeth, and on the Mode of applying it. By Mr. Robert Clarke, Surgeon at Sunderland, in the County of Durham.* 120

IX. *An Account of a new Species of Swietenia (Mahogany); and of Experiments and Observations on its Bark, made with a View to ascertain its Powers, and to compare them with those of Peruvian Bark, for which it is proposed as a Substitute:*

Being an Abstract of a Paper on this Subject, addressed to the Honourable Court of Directors of the United East-India Company. By William Roxburgh, M.D. 127

X. *An Account of the Effects of Mahogany Wood in Cases of Diarrhœa. By Mr. Francis Hughes, Surgeon of the General Infirmary at Stafford.* — — 156

XI. *Account of some Discoveries made by Mr. Galvani, of Bologna; with Experiments and Observations on them. In two Letters from Mr. Alexander Volta, F.R.S. Professor of Natural Philosophy in the University of Pavia, to Mr. Tiberius Cavallo, F.R.S.—From the Philosophical Transactions of the Royal Society of London.* — — — 162

XII. *A Return of the Sick of the Ship's Company, and of the Military, on Board the Ships in the Service of the Honourable the United East-India Company, for the Years 1792 and 1793. By John Lorimer, M.D.* — — — 211

XIII. *An Account of a singular Case of Ischuria, in a young Woman, which continued*

for

for more than three Years; during which Time, if her Urine was not drawn off with the Catheter, she frequently voided it by vomiting; and, for the last twenty Months, passed much Gravel by the Catheter, as well as by vomiting, when the Use of that Instrument was omitted, or unsuccessfully applied. To which are added some Remarks and Physiological Observations. By Isaac Senter, M. D. Associate Member of the College of Physicians of Philadelphia, and Senior Surgeon in the late American Army. Vide Transactions of the College of Physicians, of Philadelphia. — — 212

Catalogue of Books. — — 223

Index. — — 228

DIRECTIONS TO THE BINDER.

Plate the First, the references to which are explained in pages 92 and 116, may be placed at page 92; and plate the second at page 121.

MEDICAL FACTS

AND

OBSERVATIONS.

- I. *Observations on the Use of Arsenic in the Intermittent Fevers of a tropical Climate; to which is prefixed an Account of the Weather, at Sierra Leone, during the Season in which such Fevers are most prevalent. By Thomas Masterman Winterbottom, M. D. Physician to the Settlement at Sierra Leone.*

AS arsenic, though of late years frequently and successfully used in England for the cure of intermittent fevers, has not, to my knowledge, been hitherto employed in a tropical climate; some account of its use in Africa, with the histories of a few of the cases in which it was exhibited, will not, I hope, be altogether unacceptable.

VOL. VI.

B

It

It may be proper however to premise a short account of the weather at Sierra Leone during the season in which intermittents are most prevalent.

The year may be divided into the rainy, tornado, and dry seasons. The rains on this part of the coast commonly set in about the end of May, or beginning of June; and continue, more or less violently, until the beginning or middle of September: they are then succeeded by tornadoes, which continue until the end of November. It must be observed, however, that the rains are not only carried off by tornadoes, but also brought on by them; and that the tornadoes preceding the rains are, in general, less regular than those which terminate them. The dry season continues from December until May, though showers of rain sometimes occur during the dry months.

In 1792, the rains commenced about the end of May, and continued for some time to be very heavy; from the middle of July, however, until the last week of August, there were frequent intervals of fair weather, twelve hours of rain being generally followed by twenty-four or thirty hours of fair weather, with sometimes a bright sun. During this period the thermometer at noon usually stood at from 78° to 80° .

The

The last week of August and first week of September were remarkable for an almost incessant rain, which was for the most part small and drizzly, though it sometimes fell in heavy showers; the air at the same time felt cold and raw, particularly in the evenings and mornings, when a thick fog covered the hills. The thermometer at noon was from 77° to 80° .

On the 7th of September a tornado came on, which returned on the 10th, 15th, 16th, 18th, 19th, 21st, 22d, 24th, 26th, 28th, and 30th.

On the 8th, 11th, 12th, 25th, and 29th, the showers of rain were frequent.

On the 9th, 11th, 14th, and 23d, thunder and lightning occurred during some part of the day. The 9th, 13th, 15th, 17th, 24th, and 26th were sultry and almost calm. During the continuance of the rains, the winds chiefly blew from between the south and west points, but most frequently from the south-west, whence also the heaviest rain came.

As soon as the tornadoes appeared, the sea and land breezes had a more regular succession; the sea breeze usually began from the north-west about eight or nine A. M., and towards sunset drew round to the west: the land breeze then setting in from the east or south-east, con-

tinued to blow all night and during the early part of the morning.

Towards the end of the month the thermometer generally stood at 82° at noon, the atmosphere being less hazy, and the air cool.

The month of October was throughout attended with regular sea and land breezes; the atmosphere was free from haze, but sometimes overcast with clouds during the day; the whole of the month was cool and agreeable, though the thermometer at noon generally stood at 82° , and on the 29th at 84° .

A tornado occurred every night, or early in the morning, from the 1st to the 18th inclusively, frequently attended with heavy rain for some hours, and with much thunder and lightning. During the remainder of the month the tornadoes became less frequent, occurring only on the 19th, 21st, 23d, 25th, 27th, 28th, and 29th. The 1st, 17th, and 24th were sultry. On the 24th it was calm all day. On the 3d there was much thunder and lightning. On the 7th, 15th, 18th, 21st, and 30th, frequent showers of rain fell. The tornado on the 17th came from the south-west which is uncommon. The tornado on the 2d was not followed by rain. The 26th was remarkably hazy all day.

The

The lightning was extremely vivid on the 28th, appearing in long streams or chains of fire.

The month of November was much warmer than the preceding one, the thermometer at noon being from 82° to 84° . On the 11th it rose to 85° . It was on the 5th at 75° . There was continued rain till noon, when the sky became clear, the day calm and sultry. The atmosphere during the greatest part of the month was clouded and hazy, at least the tops of the hills were covered with haze during some part of the day. The sea and land breezes continued to blow very fresh, but the mornings were frequently calm and sultry till near ten A. M. On the 28th it was calm all day. Tornadoes occurred on the 2d, 8th, 10th, 12th, 13th, 16th, 19th, and 25th. The 5th, 17th, and 23d were rainy. The 5th, 11th, 14th, 18th, and 28th, were sultry, with a little wind.

In December also the sky was generally hazy and clouded; the sea and land breezes were pretty fresh during their continuance, but the mornings were for the most part calm, the sea breeze not setting in till near ten A. M.; the evenings also were close and sultry from sun-set till late at night.

A tornado came on, the morning of the 7th,

B 3 followed

followed by much rain, thunder, and lightning; but it cleared up before noon; a heavy shower fell in the afternoon of the same day.

The clearest days this month were the 3d, 9th, 13th, 18th, 24th, and 25th.

On the 5th, 8th, 14th, 15th, and 22d, gentle showers fell: on the 8th there was much thunder and lightning. The weather was sultry, with little wind, on the 1st, 3d, 14th, 19th, 22d, and 27th. The 14th and 27th were calm days. The land wind blew all day on the 13th, and the south-west and south-south west winds on the 2d, 30th, and 31st days. The thermometer at eight A. M. usually stood at from 77° to 80° ; on the 13th at 75° , and on the 26th at 81° : at noon it was from 81° to 84° ; at eight P. M. from 78° to 80° .

The remittent fever which during the months of June, July, and August, had very generally prevailed here, and had raged with great violence, began to abate in the month of September. Early in the month, this disease had not only become less frequent, but also more mild in its symptoms, gradually changing into the form of an intermittent. Towards the end of the month it became very rare, the cases which occurred being chiefly among the whites, especially

pecially those lately arrived in the country; or others who had been irregular and intemperate during the course of preceding intermittent complaints.

In the months of October, November, and December, intermittents were so prevalent, that scarcely a family in the settlement, although the whole number was nearly 400, remained perfectly free from them. They generally observed the quotidian and tertian type; there were, however, a few instances of double tertians. Most of the above cases were so mild, particularly among the men, as not to prevent them from following their different occupations, except during the time of the paroxysm. But in some instances, the daily recurrence of the disease, the long continuance of the paroxysm, and a poor diet, consisting chiefly of salted meats, rice, cassada, &c. reduced the patients to a state of great debility, and insensibly laid the foundation of long and tedious complaints. The greatest sufferers from intermittents were those who had previously laboured under remittent fevers, and had not yet recovered their strength; also persons of delicate and irritable habits, children, and women giving suck.

In every instance where the bark was taken

in due quantities, and persisted in for a proper length of time, the paroxysm was speedily checked, and the danger of a relapse effectually prevented; nor did the patient suffer those ill effects which usually occur where the disease has continued long, and been left to itself. Few, however, of the common people could be prevailed upon to take the bark in any form; and even those who took enough of it to obviate the return of a single paroxysm, would seldom continue it a sufficient length of time to eradicate the disease. These considerations, joined to an apprehension that serious and alarming consequences might ensue from frequent relapses, determined me to try the effects of the mineral solution, according to the plan recommended by Dr. Fowler*. The fear of disordering the bowels, and inducing dysenteric symptoms, rendered me at first very cautious in its use; but on finding, after repeated trials, that no ill effects were produced by its exhibition, I was encouraged to employ it more generally. The success with which it was attended will appear from the following detail of cases:

* Medical Reports of the Effects of Arsenic in the Cure of Agues, &c. 8vo. London, 1786.

CASE I.

October 4.—S. Peters, a black, aged four years, is affected every day, about noon, with coldness and violent shiverings, which continue near an hour, and are then succeeded by a hot dry skin, head-ach, and sometimes vomiting. The paroxysm is terminated in the evening by a copious perspiration. In the absence of the fit he makes no complaint, but appears languid and weak, and has little appetite. A considerable degree of hardness is felt on the left side, with a tumour projecting below the cartilages of the false ribs. He was ordered to take four drops of the mineral solution three times a day.

5. Had no cold fit yesterday at the usual time, but appeared heavy and uneasy; no sickness or griping was occasioned by the drops.

8. Has had no return of the paroxysm since the 3d. No griping nor any sensible effect has been produced by the medicine.

The solution was now omitted, and he took, on the 9th, four grains of calomel. This child had no relapse, and has continued since to enjoy

enjoy good health, although the tumour in the side did not wholly disappear till the beginning of the year 1793.

CASE II.

October 4.—Hannah Peters, a black, aged thirty-six years, has been for two months past affected with an intermittent fever; at present a paroxysm comes on every day at noon. During the hot fit, she has a considerable pain of the head, especially over the eyes, which continues till evening, and is gradually abated by the sweat which then breaks out. Her strength and appetite are much diminished.

Capiat solutionis mineralis guttas x. ter die.

6. Had no return of fever yesterday at the usual time; but towards evening had a slight cold fit, succeeded by heat and sweating. The paroxysm, however, was neither so severe, nor of so long continuance as usual. She felt a little griping in her bowels.

Repetatur Solutio.

8. Has omitted the solution two days, and has had a return of the hot fit each day at the usual time, without the preceding cold stage. She was desired to continue the drops regularly.

16. Has

16. Has taken the solution regularly since the last report, during which time she has not had the least return of her ague, nor any pain of the bowels.

Omittatur solutio et capiat Infus. Cort. Angust. ʒiij ter die.

CASE III.

October 10.—David Edmonds, a black, aged forty years, has had every day, for near a month past, a paroxysm of ague, attended with a very severe pain of the head. Of late the fit has only returned every second day, beginning about one o'clock, P.M. In the absence of the paroxysm he has no complaint but languor and debility.

Capiat solut. min. guttas x. ter die.

11. Had a slight attack yesterday evening, which did not continue long; he felt no griping or nausea from the solution.

Repetatur Solutio.

16. Has neglected his medicine for some days, during which he has missed the cold fit, but had a pretty smart hot fit every day, towards evening.

Repetatur Solutio.

20. Has had no return of the cold or hot fit
since

since the 16th: he continues the solution without experiencing any disagreeable effect from it.

CASE IV.

Octob. 5.—J. Barnes, aged thirty-six years, of a fair complexion, and florid, with red hair, was attacked with the remittent fever about the end of August last, from which he recovered by a liberal use of the bark; but soon after, on returning to work, and exposing himself too much in the sun, he suffered a severe relapse in the beginning of September. His complaint, however, yielded again to the bark, but left him greatly enfeebled. During the remainder of the month of September, he continued to take from $\mathfrak{z}\text{i}$ to $\mathfrak{z}\text{i}\mathfrak{s}$ of bark every day, and returned to his work. About a week afterwards he was suddenly seized with a cold fit, followed by a hot stage and a profuse perspiration, which left him very weak during the apyrexia. His pulse is now 100, rather hard and quick: he has a severe attack every day at noon, attended with vomiting, and, during the hot fit, with a quick and hurried respiration; he is hot
and

and restless till late in the evening, and has then very profuse night sweats.

Capiat solut. min. guttas x. ter die.

Oct. 6. The solution did not disagree with him. He had a slight return of the paroxysm yesterday.

Repétatur Solutio.

8. Has had no return of the fit, nor felt any sensible effect from the medicine. He perspired much at night; has great debility and languor, with little appetite.

10. The symptoms are nearly as before; he did not rest well, but had no return of the paroxysm.

Capiat opii gr. ij h. s. Repetatur Solutio.

12 The solution was yesterday omitted; he rested better with the pill: in other respects finds no alteration.

13. Had a return of the paroxysm yesterday; the cold stage lasted half an hour, the hot stage about two hours. He was much relieved by the opium, and sweated very profusely after it.

14. Had another slight fit yesterday evening, the cold stage being very short; he sweated much: does not recover his strength or appetite. As he could not be prevailed upon to take the bark again, I directed that four ounces of the following

following infusion should be taken three times a day:

R. Corticis Angusturæ ʒj Cremor. Tart. ʒij Aquæ pur. ℥iiss.

By this plan his appetite became better, and he regained his strength in some degree; but in a week or ten days he relapsed into his former state, having every day an ague fit, which was, however, relieved by two grains of opium, taken at the commencement of the cold stage. He now began to take the bark to the amount of ʒiiss a day, which finally put a stop to the ague; notwithstanding, he recovered his strength so slowly, that it was thought necessary, six weeks afterward, to send him to England for the effectual restoration of his health.

CASE V.

October 14.—A. Richardson, a black, aged forty years, since her recovery from a remittent fever in August last, has continued in a very debilitated state, and for some time past has been affected with an intermittent fever, the cold fit of which comes on daily at four o'clock, P. M. is very severe, and of long duration.
Much

Much pain of the head, and frequent vomiting attend the hot fit, which continues the greatest part of the night, and is succeeded towards morning by a slight partial sweat: she remains very weak till the commencement of the next paroxysm; her appetite is much impaired; her body open.

Capiat solut. min. guttas x. ter die.

16. Has taken the solution two days, and has had no appearance of the ague, except a little uneasiness and yawning about the time of its usual attack. No sensible effect is produced by the medicine.

Repetatur Solutio.

17. Had a return of the paroxysm yesterday; the cold fit was short, but severe; the hot fit was also violent, and terminated by a profuse perspiration; after which, however, she appeared more easy and composed than usual. She complained of no griping or nausea from the medicine.

Repetatur Solutio.

24. Has had no return of the paroxysm since the 17th, nor any symptoms of its approach. She continues still very weak, and has little appetite.

Omittatur solut. Capiat infus. gent. c. ʒij ter die.

28. Has had no return of the fit. She begins

gins to recover her strength and appetite.

Repetatur Infus.

CASE VI.

Nov. 2.—Mary Bowler, aged forty years, a black, has been for six weeks affected with a tertian ague; the cold fit is severe; the hot fit, which is very violent, and attended with great pain of the head, generally continues all night, and sometimes part of the next day, without any sweating stage. She is much debilitated, but has a tolerable appetite.

Capiat solut. min. guttas x. ter die.

4. Had a return of the paroxysm yesterday, after the third dose of the solution. The fit returned at the usual period, and in the same manner as before. No sensible effect was produced by the medicine.

Repetatur Solutio.

8. Has had no return of the cold fit since the 3d; the hot fit occurred about the usual time, but it was shorter and much less severe than ordinary.

Repetatur Solutio.

12. Has had no return of the paroxysm; she complains of a little griping in her bowels, and continues still weak.

Omittatur solut. min. Capiat inf. gent. c. ʒij ter die.

20. She makes no complaint, and has nearly recovered her health and spirits.

Repetatur Infus. Gent. c.

CASE VII.

Nov. 1.—E. Perth, a black, aged forty-five years, has been for near six weeks past affected with an irregular intermittent, which most commonly follows the tertian type. The cold fit is severe, and very uncertain in the time of its attack and in its duration. In the hot fit she complains of excessive pain of the head, especially over her eyes, and of great pain of the back. The hot stage generally continues all night, seldom terminating by regular sweats: it is followed by much lassitude and uneasiness through the ensuing day. Her strength is greatly impaired, her appetite bad; and she is very costive.

*Capiat statim Sal. cathart. amar. ʒi. Cras incipiat sumere
Sol. min. guttas x. ter die.*

6. After taking three doses of the medicine, she had a return of the paroxysm on the 3d, but thought the cold fit later in its approach than usual, and shorter. The hot fit continued

VOL. VI.

C

throughⁿ

through a great part of the night, but the pain of the head was much less severe. She has had no return of the paroxysm since, and feels only a little griping from the medicine.

☞ Repetatur Solutio.

10. Has had no return of the paroxysm since the 3d. She complains only of debility and want of appetite.

Omitatur Solut. Capiat Infus. Gent. c. ʒij ter die.

14. Begins to recover her strength; her appetite is also better.

Repetatur Infus. Gent. c.

CASE VIII.

Octob. 3.—Ann Bowler, a black, aged fourteen years, has been, for some weeks past, affected with an irregular tertian, which is sometimes, but not generally, preceded by a cold stage. The hot stage continues during the greater part of the day, and seldom terminates by sweating. Her body is open; her appetite much impaired.

Capiat Solut. min. guttas viii; ter die.

10. The solution has now been taken for a week, during which time she has had no return
of

of the ague, nor has felt any nausea or griping from the medicine. No complaint remains but debility.

CASE IX.

October 4.—Dinah Lawrence, a black, aged forty-four years, is every other day, about six o'clock P. M., seized with a severe cold fit, followed by great heat and violent pain of the head, especially over the eyes, which symptoms continue through the whole night, and are not succeeded by any regular sweating stage; she is costive, and much debilitated; she has had this complaint near three months.

Capiat statim Sal. cath. am. ʒvi; et cras Solut. min. guttas x. ter die.

10. Has had no return of the fit since she began to take the solution; she finds no disagreeable effect from it: is still costive.

Repetantur Sal cathart. et Solut. min. ut antea.

14. Feels no complaint but what proceeds from debility; her appetite is better; she was a little griped by the medicine.

Omittatur Solut. min. Capiat Infus. Gent. c. ʒij ter die.

CASE X.

Sept. 24.—Jane Armstrong, of a fair complexion, aged thirty years, is seized every day, at eleven o'clock A. M., with a head-ach so violent as to produce frequent shrieking and continual moaning. The pain chiefly affects the crown and one side of the head; it is in general preceded by a cold stage, though slight, and of short duration. The hot fit, which is not very violent, continues till night, when it abates along with the pain; but is not entirely removed till morning: the paroxysm is usually terminated by a profuse perspiration. The patient is naturally of a delicate constitution, and has of late been much reduced by the remittent fever, from which she recovered very slowly.

Capiat Opii gr. iij et Tart. emet. gr. $\frac{1}{2}$ ingruente paroxysmo.

25. The head-ach was almost entirely removed within half an hour after taking the pill; the paroxysm terminated also more speedily than usual. Being very costive, she was ordered to take half an ounce of purging salt the following morning.

26. The

26. The salt operated gently; she had a very violent return of head-ach at the usual time, which was relieved by the opium taken alone.

October 4.—She refuses to take the bark: she has every day had a return of head-ach at the usual time, which was however removed by the opium.

Capiat Solut. min. guttas x. ter die; et repetatur Opium sub initium paroxyfini.

10. Has had no return of the paroxysm since she began the solution; feels no inconvenience from its use, but a slight diarrhoea, without any pain.

Repetatur Solutio.

14. Has had no return of the head-ach; she sweats much at night; is very weak, and has no appetite.

Omittatur Solut. Capiat Infus. Cort. Angust. ℥iij ter die.

This woman has never had a return of the paroxysm, though a twelvemonth has now elapsed since the last report. She gradually recovered her strength by the use of tonic remedies.

CASE XI.

Sept. 12.—Jesse George, a black, aged twenty
C 3 years,

years, was yesterday afternoon seized with a severe cold fit of an ague, which continued upwards of two hours, and was succeeded by great heat, severe pain of the head, nausea, pains all over his body, more especially in the back and loins, great restlessness, and anxiety. Towards morning a general but not profuse perspiration took place; the severity of the head-ach at the same time abated, and all the other symptoms wholly disappeared: he has much thirst; his skin is cool; his pulse 72, and soft.

Capiat Solut. min. guttas x. ter die.

13. He had a return of the paroxysm last night, at eight o'clock, four hours later than the former one. The cold fit, though very severe, did not continue long; the hot fit was strong; the head-ach less violent. He had a very profuse perspiration this morning. His skin is now cool and moist, and his tongue clean; but some pain still remains over the orbits of the eyes; he complains of thirst, and is costive.

Capiat Sal. cathart. am. ʒi—Repetatur Solut. min.

14. The head-ach continued yesterday till the afternoon, and then went off; the salts were not taken till this morning. He rested well last night, and makes no complaint but of debility.

Repetatur Solutio.

15. He

15. He continued free from complaint yesterday, till towards evening, when he became hot and feverish; and after a very uneasy night, he, this morning, at eight o'clock, had a severe cold fit, attended with violent head-ach, which lasted near an hour. Two grains of opium, taken at this time, brought on a sweat, and terminated the paroxysm.

Repetatur Solutio.

16. He slept well last night, and feels no complaint but from debility. He has omitted the drops this day.

Repetatur cras Solutio.

17. Has had no return of the paroxysm; he feels no complaint but a slight griping from the solution.

R. Tinct. Opii et Solut. min. aa ʒij m. capiat guttas xx. ter die.

20. He has had no return of the paroxysm since the 15th. At that time he probably brought it on by having exposed himself the night before to the damp evening air in his shirt. He feels no griping, or sickness, from the drops, which he still takes. He returned to his work this day.

CASE XII.

August 12, 1793.—Mr. T——, a European, of a dark complexion, with black hair, was suddenly seized, two days ago, with an acute pain of the head, chiefly over the orbits of the eyes, attended with nausea and vomiting. These symptoms were soon followed by great heat and restlessness, which continued through the whole night, and yielded in the morning to a profuse perspiration. On the 11th he was free from complaint; walked about, and ate heartily. In the evening, however, he was seized with a very severe shivering fit, which continued near two hours, and was succeeded by great heat and restlessness, by severe pain above the eyes, and bilious vomiting. He was again relieved in the morning by a copious perspiration. At ten o'clock, A.M. his skin was still hotter than natural, and his pulse rather quick; in other respects he appeared free from complaint.

Capiat Solut. min. guttas x. ter die.

13. The first dose of the solution yesterday produced vomiting; the second gave him three stools;

stools; the last had no particular effect. He passed an easy night, without feeling any symptom of the fit, except a general uneasiness, which, however, soon went off. He complains this morning of slight pain over his forehead.

Repetatur Solutio.

14. The medicine again produced sickness, and a slight diarrhœa, though he only took two doses of it. He remained well till two o'clock, P.M.; he then became very hot, and had a severe return of the head-ach, attended with nausea and vomiting. The heat, pain, and restlessness continued till this morning, when a copious perspiration took place, with which he is yet affected.

At ten o'clock A.M. his pulse is 130; his skin pretty cool; his head-ach almost gone; his tongue somewhat furred. He complains of thirst, and of slight pain of his bowels, with a sensation of numbness about the umbilicus

Omittatur Solut. Capiat pulv. Cort. Peruv. ʒi secunda quaque hora.

At six o'clock, P.M. he has a very slight head-ach, with a sense of weight in the forehead; his eyes are more prominent and brighter than usual. He has taken two doses of bark since noon, the first of which produced vomiting; he has
had

had one stool to-day; his urine is very high coloured; pulse 130, soft, and less quick than in the morning.

Repetatur Cort. et capiat h. s. Tinct. Opii et Vin. Antim. aa guttas xxx.

15th. Ten o'clock, A.M—he has had a good night; some pain still remains over his eyes, but it is less severe; his skin is rather hot, but moist; pulse 112; his tongue dry and white; his urine high coloured, with a light cloud suspended in it. He complains much of thirst and fever, and of a pain in his back. He has taken, since yesterday noon, ʒiss of Peruvian bark.

Repetantur Cortex, Tinct. Opii, et Vin. Antim.

16. He passed an easy night, and enjoyed some refreshing sleep; he complains only of a slight pain over his eyes, and is able to sit up. He had two stools in the night; his tongue is cleaner, but still dry; pulse 104 and soft, but easily quickened by the least exertion. His urine is not so high coloured, and exhibits a flocculent cloud. He took ʒiss of bark between ten o'clock, A. M. yesterday, and six o'clock this morning.

Reperantur Cortex, Tinct. Opii, et Vin. Antim.

17. He was much griped yesterday by drinking some cyder; has no complaint this morning

ing but from weakness. His pulse is 104, and soft; his tongue clean and moist. His urine is much paler than before, and has a kind of gelatinous striated cloud suspended in it.

The same medicines were repeated.

18. He seems much better in every respect; his appetite is returning; his pulse 90, and soft.

He continued the bark a few days longer, and had no return of complaint.

CASE XIII.

October 4.—Ann and Eliz. Davis, blacks, the former five, the latter six years old, have been for some time past affected with quotidian agues. The cold fit comes on at four o'clock, P. M.; is very severe, and frequently attended with vomiting. The hot fit usually continues the whole night, being attended with great restlessness, anxiety, and acute pain over the eyes; but is seldom succeeded by a regular sweating stage. Their appetite and strength are much impaired.

Capiant Solut. min. guttas vj. ter die.

9. Each of them had a return of the cold fit
on

on the 4th, after the third dose of the solution. They have since had no return.

Repetatur Solutio.

11. There has not been any appearance of the paroxysm, nor any disagreeable effect from the medicine.

CASE XIV.

John Oliver, a black, aged five years, who was affected nearly in the same manner as the two last patients, began, on the 16th of August, to take four drops of the solution three times a day.

23. He had a return of the fit on the 16th, 17th, and 18th, but it commenced every day later, was less severe, and of shorter duration. Since the 18th he has had no fit, although the solution was discontinued. A slight tumefaction of the face has been observed for two days past, but is at present subsiding. He felt no nausea or pain from the medicine.

CASE

CASE XV.

Dec. 10.—Mary Jones, a black, aged thirty-six years, about three months ago was affected with a remittent fever, from which she recovered very slowly, and has since continued in a state of great debility. She has of late been subject to violent pains in the bowels, attended with diarrhœa. During the last month she has had a regular tertian ague, the cold fit of which begins generally at sun-set, but is not very severe, nor of long continuance. The hot fit is long and severe, being attended with violent head-ach, intense thirst, and great restlessness. These symptoms are not terminated by a regular sweating stage; and have often no remission till the middle of the following day. She is feeble, and much emaciated.

Capiat Solut. min. guttas x. ter die; et Opii. gr. ij, sub accessionem paroxysmi.

12. The hot fit was much relieved by the opium; the paroxysm was shorter, and the head-ach less severe. She is very costive.

Repetatur Solut. min. et capiat Sal. cathart. ℥ss mane.

15. Con-

15. Continues the solution without feeling any sensible effect from it. She has had no cold fit or head-ach during the two last paroxysms. The hot fit was much less violent and of shorter duration than formerly.

Repetatur Solutio.

18. Has had no return of the fit, nor any appearance of it since the last report; nor does she perceive any nausea or griping from the solution. Her appetite is still much impaired.

Repetatur Solutio. Capiat Inf. Cort. Angust. ʒij ter die.

22. There has been no return of the paroxysm. She finds her strength and appetite much increased by the infusion.

The use of the solution was discontinued.

CASE XVI.

Feb. 1.—John Jones, a European, of a fallow complexion, aged twenty-eight years, is affected in the afternoon, every other day, with a violent cold fit, attended with rigors, and succeeded by a regular hot fit and sweating. Until within a few days, he has been able to do his duty on ship-board as a seaman; but the paroxysm returns now with so much violence,

as to confine him to his hammock. He has taken a large quantity of Peruvian bark at different times, which has never failed to prevent the next return of the paroxysm; he has always, however, had a relapse in a few days, through intemperance, and exposure to the night air.

Capiat Solut. min. guttas x. ter die.

8. Has taken the solution without perceiving any sensible effect from it. The paroxysm returns as usual, but, as he says, with much less violence.

Repetatur Solutio.

15. The paroxysm returns as usual, but is shorter and less severe. Through mistake, he has taken the solution only before the attack of each paroxysm.

Repetatur Solutio; et capiat guttas x. ter die.

20. He has had no return of the paroxysm since he took the solution as directed, and feels no nausea or griping from it.

He continued the medicine a few days longer, and was restored to perfect health.

CASE XVII.

Feb. 1.—Ann Wicks, a mulatto, aged forty years, has been for a month past affected, every other day, with a violent cold fit, attended with rigors, and succeeded by great heat. She has also a severe pain over the forehead, and on one side of the head, extending to the neck and shoulder of the same side. There is much stiffness and pain in moving the neck during the intermission. The cold stage commences about five o'clock, P. M. and continues near an hour. The hot fit does not terminate before morning, and is seldom succeeded by a regular sweating stage. She is much debilitated by the long continuance of the complaint, and has lately given suck to a young child. Her appetite is also greatly impaired.

Capiat Solut. min. guttas viij ter die.

4. She has had no return of the cold fit. The hot fit continued only part of the night, and was unattended with head-ach or any other distressing symptom.

Repetatur Solutio.

12. She

12. She has had no return of the paroxysm, and feels no ill effects from the solution. Her strength is somewhat increased, but her appetite is still bad.

Omittatur Solutio. Capiat Inf. Gent. c. ζ iss bis die,

CASE XVIII.

Mrs. D. a delicate woman, of a fair complexion, aged twenty-four years, in the month of August last had a miscarriage, from which she recovered without much trouble, and enjoyed a tolerable state of health till the beginning of October, when she was seized with the common remittent fever of the place. From this complaint she also recovered within a fortnight, by taking largely of the bark in powder and decoction. About the end of the month, however, she suffered a relapse, and made a very slow progress towards recovery; her stomach being only able to retain the bark in the form of a decoction. She laboured under great debility, very profuse night sweats, and frequent hectic flushings during the day, with loss of appetite, and general tremors on using the least exercise. These symptoms were at length

Vol. VI.

D

confide-

considerably alleviated by the infusion of Angustura bark, elixir of vitriol, and other tonics.

Dec. 15. Yesterday, at six o'clock, P.M. she had a cold fit, with rigors, which lasted near half an hour, and was succeeded by a hot fit, attended with great pain of the head, nausea, vomiting, and restlessness, which continued through the whole night; towards morning she was relieved by a partial sweat, but remained very weak and languid.

16. Yesterday, at the same hour, she had a return of the paroxysm, the symptoms of which were mitigated by an opiate taken soon after its commencement: she had a copious perspiration during the night, and seems free from complaint this morning.

18. Had a return of the fit on the 16th and 17th, but was relieved as before by an opiate. She refuses to take bark.

Capiat Solut. min. guttas viij. ter die.

20. She has had no return of the cold fit since the 18th. The hot fit was much shorter and less severe. She experiences no inconvenience from the medicine.

Repetatur Solutio.

22. She has had no return of the paroxysm, but feels a slight pain in her bowels.

Capiat statim Tinct. Opii guttas xx. et sp. lav. c. 3ss.

Repetatur Solut. min.

24. The

24. The pain of her bowels was removed by the opiate; she has had no return of the paroxysm; rests pretty well during the night, but sweats much towards morning.

Omittatur Solutio; et capiat Infus. Gent. c. ʒi ter die.

30. Her strength is returning. Her appetite is good, and she has had no return of the paroxysm.

This lady continued to enjoy a good state of health, till the 20th of March, 1793, when she was affected with a diarrhœa, attended with acute pain in her bowels, chiefly about the umbilicus. She was soon relieved from these complaints by an opiate, and a few powders, consisting of the colombo root joined to an aromatic: but on the 25th, she had a return of an intermittent fever, the cold fit of which was very severe. It began at six o'clock in the evening, continued near two hours, and was followed by a hot fit, which lasted all night, terminating towards the morning in a slight perspiration, and leaving her low and weak the remainder of the day.

28. She refused yesterday to take an opiate on the approach of the cold fit, having on former occasions found her head disagreeably affected by it. The paroxysm proved very severe: the

hot fit continued all night, and was succeeded by partial sweats about the head and neck. She is very weak this morning, and complains of a great pain of the head and back; of lowness of spirits and general uneasiness.

Capiat Solut. min. guttas viij. ter die, ex Infus. Cort. Angustur. cyatho.

30. The solution did not disagree with her in any respect; she had a cold fit last night, but it was much less severe than usual; she is also in better spirits to-day.

Repetatur Solutio.

April 1. There was no cold fit yesterday; but she had a hot fit, which continued all night, and terminated in a very profuse perspiration. Her spirits are much revived; she is considerably stronger, and has a better appetite.

Repetatur Solutio.

6. She continues the solution without feeling any inconvenience from it; and has had no return of the fit, or night-sweats, since the 1st: her appetite at present is good.

Repetatur Solutio.

8. She has had no fit, and recovers her strength gradually. No nausea or griping has ever been produced by the solution.

Omittatur Solutio. Capiat pulv. rad. colomb. gr. xv. ter die.

CASE

CASE XIX.

Feb. 1, 1793—Mrs. H. of a fair complexion, aged twenty-four years, during the months of September and October last, had two several attacks of the remittent fever, from which she recovered speedily by means of the bark : since that time she has continued in a very weak irritable state, subject to pains of the bowels, and to frequent though slight returns of a febrile complaint, which continued only for a day or two, and commonly yielded to an opiate. On the 27th of January she had a cold fit at eight o'clock in the morning ; this was succeeded, in about an hour, by a burning heat of the skin, with flushing of the face, great restlessness, and severe pain of the forehead. Her eyes, at the same time, appeared bright and prominent ; she complained also of a sense of heat in them, and was unable to bear the light. In the evening, a copious perspiration ensued, and considerably alleviated the symptoms ; she had, however, a slight head-ach through the whole night : the

fit has returned every morning at the same time for the last four days.

Feb. 2. The paroxysm appeared this morning as usual, with a severe cold fit and head-ach, but was rendered much shorter and less distressing by an opiate draught taken soon after its accession.

Capiat Solut. min. guttas viij. ter dic.

5. She had a slight return of the cold fit this morning, with a little head-ach, but the paroxysm was of short duration.

Repetatur Solutio.

6. She has had no cold fit to-day, nor any pain of the head; the hot fit returned at the usual time. Her face is much flushed, and her skin hot, but with less anxiety and restlessness than heretofore: she finds no inconvenience from the solution. The opiate was not taken to-day.

Repetatur Solutio.

10. She has had no return of the paroxysm, nor has felt the slightest symptom of its approach since the 6th; she complains only of a slight pain or uneasiness in her stomach. Her appetite still continues weak.

Omittatur Solut. min. Capiat tinct. opii guttas xx. statim.

14. She begins to recover her strength and appetite;

appetite ; the pain of the stomach was immediately removed by the opiate.

All the patients whose cases are here related, have continued to enjoy good health since cured by the solution; and though several months have now elapsed, none of them have experienced the least unpleasant symptom which could be attributed to that remedy. The women continued to labour under a suppression of the catamenia, until their strength was entirely restored.

Mrs. H. (Case XIX.) though enjoying a good state of health, had no appearance of them till the middle of August last, when they flowed for several days rather profusely.

In Case IV. I had little prospect of success from the use of the solution, the child having become very weak and irritable by frequent relapses : but as he had for a length of time taken the bark in large doses without any effect, I was induced to try the mineral solution, with a view of checking the returns of the paroxysm, hoping afterwards to complete the cure by the bark ; which might prove more effectual after its use had been suspended a few days.

In Cases I. X. XIII. and XIV. there was an evident enlargement of the spleen, forming a projection below the cartilages of the ribs. In Case X. it was so large as to extend nearly as low as the crista of the os ilium. After the ague had ceased, the patient continued to use corroborant medicines, taking at the same time small doses of calomel, but without any sensible effect on the tumor; it yet remains nearly in the same state, not, however, causing much uneasiness. In Cases XIII. and XIV. as the patients speedily regained their health after the ague had ceased, and felt no uneasiness from the enlargement of the spleen, I did not think it proper to use any medicine, excepting a purgative dose of calomel occasionally, because, in many similar cases, where this medicine had been used, even in very small doses, a salivation was very soon excited, the tumor not being at all affected by it, whereas the patient was rendered extremely weak and irritable. The only instance of tumefaction which could with any probability be referred to the use of the solution, was Case XIV. in which, however, it proved so slight, as scarcely to deserve notice.

In order to give the mineral solution a fairer trial, I avoided, in many instances, making
use

use of two very powerful means usually employed for the purpose of diminishing the violence of the paroxysm, and which frequently indeed put a total stop to it; I mean, opium and emetics: when two grains of opium are given a short time before the paroxysm is expected, it seldom fails to bring the fit to a speedy termination by a profuse sweat; and generally relieves the violent pain of the head, which is so distressing during the hot fit, as in Cases X. and XV. The recurrence of the paroxysm being once obviated, I have found that a full dose of opium at night affords more comfortable rest, and more certainly prevents the solution from affecting the bowels, than when the tincture of it is added to the mineral solution; a mixture of this kind always becomes turbid, and the opium is partly separated.

Intermittents partake much of the nature of remittents, and the two diseases have a very uncertain boundary; whenever, therefore, the intermissions are imperfect and indistinct, the exhibition of an emetic is attended with most beneficial effects. In many instances this practice puts a temporary stop to the returns of the fit, and in every case considerably diminishes its violence. The proper time of giving an
 emetic

emetic, is about two hours before the paroxysm is expected; and the best mode is to employ a solution of tartarized antimony in divided doses, at intervals of eight or ten minutes, until full vomiting be produced. When the patient has vomited a few times, and his stomach is a little settled, a more moderate dose of the antimonial solution, joined to a full dose of opium, seldom fails to produce a copious perspiration before the attack of the cold fit. This method generally succeeds in preventing the immediate recurrence of the paroxysm: but in those cases where the intermittent has continued long, and seems to return by the power of habit, it will be proper to repeat the emetic once or twice more before the time when the paroxysms are expected.

I think it proper here to observe, that antimonials, in the nauseating doses in which they are frequently given during the remission or apyrexia, with a view of procuring a more perfect solution of the disease, are seldom found adequate to the purpose; on the contrary, the continued action of so powerful a stimulus, in general, produces a correspondent state of debility, and relaxes the muscular fibres of the stomach

mach so much, that neither food nor medicine can be properly retained.

The remittent fever is, in many cases, very mild; whence the remission has often been mistaken for an intermission. This mistake is more liable to be made when the remittent fever is preceded by an evident and severe cold stage at each return of the paroxysm, and is followed by a regular hot, and sweating stage. The remittent may, however, be distinguished from the intermittent fever; 1st, by a slight pain which remains fixed in the forehead, or over the orbits of the eyes, during the apyrexia; 2dly, by the pulse, which, though not more frequent than in health, yet retains a degree of quickness or sharpness through the whole of the remission; 3dly, by the state of the skin, which, though moist, feels hotter than natural. In such cases I have not found the mineral solution so successful as in those where the intermission was complete; for which reason it seems most prudent to place our sole dependance upon the bark, as in Cases IV. and XII. Sometimes, however, when the patient could not be prevailed upon to take the bark in proper doses, I have found much advantage from joining it with the mineral solution, by which means a smaller quantity

tity of bark will answer the intended purpose. But whenever immediate danger presents itself, or is to be apprehended from a continuance of the fever, the bark, given in large doses, is the only medicine to be depended on.

The mineral solution usually fails in some irregular cases, which at first view resemble intermittents, and have been improperly ranked with them, under the denomination of erratic or anomalous intermittents. A morbid increase of irritability appears to be the foundation of these irregular complaints; they affect principally those who have been debilitated by frequent attacks of fever, or by lingering diseases; also children; and women, more especially those who give suck; and, in general, persons of a weak delicate habit. The symptoms which occur in these complaints are nearly as follow: during the afternoon, or towards evening, the patient becomes uneasy and restless; his skin feels dry, and is hotter than usual, but without imparting the burning heat usually observed in the hot stage of intermittents; the pulse becomes quick, and rather more frequent than natural; a pain is sometimes felt in the head, either on the crown, or on the back part of it; the thirst is seldom very great; disagreeable
clammi-

clamminess, however, takes place in the morning. These symptoms are sometimes preceded by slight chills running down the back, which, however, when they do occur, are not of long continuance, and never accompanied with violent shiverings.

In this manner the patient is harrassed during the whole night*, but obtains relief towards morning, when a partial sweat sometimes appears about the head and breast. Excepting a degree of languor and debility, little or no complaint is felt till the return of evening. The duration of these complaints is very uncertain; they sometimes affect the patient daily for one or more weeks; at other times abate or disappear for a few days, and then return as before. Whatever increases the irritability of the body, may be considered as an occasional cause of them; but the most common as well as most powerful one is too much fatigue, along with exposure to a hot sun.

In these cases, after evacuating the stomach and bowels by a gentle emetic or purgative, it is commonly sufficient to exhibit some tonic, in a form agreeable to the patient's stomach. The

* Hence the denomination of night-fever.

Peruvian bark does not appear to produce any better effects than the other vegetable tonics, as Gentian, Colombo, &c. An infusion of Angustura bark is what I most frequently employ, and find most useful, taking care to prevent the costiveness arising from its use, by giving, at proper intervals, a dose of calomel.

For children, who cannot easily be induced to take bitters, after the previous use of an emetic, a few moderate doses of calomel are commonly sufficient.

Notwithstanding the effects of arsenic appear to be equally as powerful and nearly as certain as those of bark in the cure of intermittent fevers, yet it must be confessed that perfect strength is less speedily recovered when the cure has been accomplished by arsenic alone, than when bark has been employed. This objection to the use of arsenic is of less consequence in cold climates, where, if the ague has not been of long standing, the debility induced by it is seldom very considerable. In tropical countries, however, a few attacks of an intermittent frequently reduce the patient so much, that even when the paroxysm has ceased to return, the extreme debility which remains, is
of

of itself sufficiently alarming to demand every attention from the practitioner.

It does not appear improbable that the bark owes its specific power, in the cure of remittent and intermittent fevers, to some peculiar principle in its composition, which has hitherto eluded the researches of experimenters, and which they have in vain attempted to imitate by various combinations of bitters and astringents. In whatever this peculiar power of the bark may consist, the same quality appears to be possessed by the arsenic in a considerable degree. Both remedies probably effect the cure of intermittents, by their action upon the fibres of the stomach, since they often operate speedily, and even in a small dose; but the power of the arsenic seems to cease here; whereas the bark is capable of restoring tone to the system in general. The same effect may perhaps be nearly obtained by joining some tonic medicine to the arsenic. With this view, in many cases, after the solution had been taken a week or ten days, I discontinued its use, and ordered the patients to take the Infus. Angust. Infus. Gent. c. &c. until their strength was completely restored. It may be found still more advantageous to employ

ploy these remedies along with the mineral solution.

Arsenic seems to have been oftener employed as a medicine in Germany, than in any other part of Europe ; but chiefly by the empirical class of practitioners, which no doubt prevented its introduction into general use. Many eminent physicians in Germany, as well as elsewhere, have, however, spoken highly in its favour, and occasionally prescribed it. Like many other active remedies, it has been much abused by the bold and the ignorant, and has been given in doses which no man of prudence would venture to direct ; especially as we know that the same good effects may be obtained by moderate doses of it, and without the least risk. The following observations, extracted from a German work *, will show how extensively this medicine has been used on the Continent, and how little caution has been observed in its exhibition.

Dr. Slevogt, Professor of Anatomy at Jena, in 1700, recommended the use of arsenic, extolling it as the best, most certain, and safest cure of intermittents, especially of tertians and quartans. He employed it in doses of a grain

* Nicolai Recepten und Kurarten. 8vo. Jena, 1780.

of a grain and a half mixed with a proper quantity of Theriaca; not only giving it on the days of the apyrexia; but also a short time before the accession of each paroxysm. He asserts, that in fifty instances, two or three doses were sufficient to put a total stop to the disease, and that he never observed the least ill effect from it.

Melchior Frickius* recommends arsenic in intermittents, and declares he has used many drachms of it in the cure of such fevers; but confesses that he had often met with relapses afterwards.

Lanzonus † quotes a letter from Valisnieri to one of his friends, written in 1707, in which he says the French surgeons were accustomed to cure long-continued intermittents with a small quantity of arsenic: and he adds, that their remedy seemed to resemble much the famous aqua del petefino, which was a strong solution of arsenic boiled in a copper vessel ‡.

* De Virtute Venenorum medica. 8vo. Ulmæ, 1701.—
See also London Medical Journal, Vol. VII. p. 194.

† Lanzoni Oper. omn. med. phys. 4to. Lauf. 1738.
Tom. I. p. 68.

‡ The *Aqua della Toffanina* (so called from the inventor), *Aquetta di Napoli*, *Poudre de Succession*, *Eau Mirable*, &c. were preparations of arsenic frequently used as poisons during the last century.

Keil * praises arsenic as a certain and safe specific in intermittents, when prepared and administered in the following manner: half an ounce of white arsenic, finely powdered, is to be put into a glass, or tea-cup; half an ounce of distilled vinegar is then to be added, and evaporated over the fire, being constantly stirred at the same time with a wooden spatula; the same quantity of vinegar is again to be added and evaporated in like manner. After this process has been repeated six times, the residuum is finally to be washed with warm water, and dried; a drachm of the dry powder is to be made up into sixty pills by means of a scruple of wafers softened with water. Previously to the use of the pills, the patient is to take an emetic composed of tart. emet. or sulph. aurat. antim. and a little vitriolated tartar, or some purgative medicine on the morning free from fever: the next day, or only a few hours before the accession of the paroxysm, one of the pills is to be taken fasting, and nothing is to be eaten or drank after it for three or four hours. When this has been repeated three days, during the apyrexia, or a few hours before the

* Anatom. Chirurg. Medicin. Handbuchlein. 8vo. Koenigsberg, 1761.

attack of the paroxysm, the fever commonly ceases. He affirms that this practice has been attended with success in several hundred cases, when every other remedy had been employed in vain; that he has never observed the least ill effect to accrue from it; but, on the contrary, that those who had before looked thin and ill, had become, in consequence of it, fat and strong; and that he knew many persons who had used this remedy fifteen or twenty years before, and who continued to enjoy a state of perfect health.

Dr. Jacobi * recommends the use of arsenic strongly in fevers: he directs one part of arsenic and twelve of salt of tartar, to be mixed with 180 parts of water, and boiled till one half has evaporated; when cold, as much fresh water is to be added to it as has been lost by the evaporation, together with a little spirit of wine. The dose for adults is twenty-five drops, to be given on the day which is free from fever, at seven A. M., at three, six, and nine, P. M. Before the use of this medicine, the *primæ viæ* must be evacuated by emetics and purgatives; and the common febrile

* De prudenti usu Arsenici, sale Alcalico domiti, interno salutari, Differt.—Vide Act. Acad. Elect. Mogunt. Tom. I. p. 216. 8vo. Erford. 1751.

fuge remedies should be used for some time. Dr. Jacobi observes that he has employed the above preparation not only in intermittents, but also in continued fevers, with the greatest success, and without ever experiencing any bad effects from it.

Heuermann * says that arsenic is used in Holstein, at Copenhagen, and some other places, as the most certain remedy for the cure of intermittents; that he has himself given it with constant success, in fevers, to patients who were not able to retain other medicines on the stomach in a proper quantity; and that two cases, wherein frequent relapses had occurred, were entirely cured by this remedy. He prepares a solution of arsenic in the following manner: half an ounce of white arsenic, and six ounces of alkaline salt, are added to lbs of water, and then evaporated to dryness. The same quantity of water is added a second time to the residuum, and evaporated to one half, which is coloured red by a few poppies. Of this he directs from seven to ten drops to be taken during the day, beginning immediately after the paroxysm is over, and omitting it a short time before the return of the next. If the

* Vermischte Bemerkungen und Untersuchungen. Vol. I. 8vo. Copenhagen, 1765.

solution produces vomiting, it is too strong, and must be diluted; only one dose is to be given in twenty-four hours, and the patient must be kept moderately warm, to promote a gentle perspiration. Exposure to cold, he says, is as hurtful during the use of this as of other febrifuge remedies, as it disturbs Nature in her operations, and retains in the body the noxious matters which she is endeavouring to expel. If in the first three or four days after the use of these drops, the fever does not cease, he recommends that the same dose should be repeated twice a day, which commonly proves sufficient. The ill consequences which have been observed after the use of arsenic, as palsy, trembling of the limbs, blindness, deafness, &c. he ascribes to the improper preparation and imprudent use of it; asserting, that it is a safe remedy when properly prepared.

In the Ephemerid. Acad. Nat. Curios.* arsenic is also celebrated as an infallible specific for intermittents. Three or four grains of powdered white arsenic are directed to be put into a small uncovered glass with a proper quantity of water, and placed upon the fire till a solution takes place, when it is to be well stirred up and given to the patient: the fever, we are in-

* Dec. II. Ann. III. p. 132.

formed, is by this means certainly prevented from returning. The patient should eat nothing for twelve hours before; but a quarter of an hour after having taken the medicine, he is allowed a gill of warm water, in which a quantity of butter is dissolved, together with the yolk of an egg; after which, nothing more is to be given for some hours. There generally follows a considerable degree of uneasiness, and a profuse sweat; and by these means, it is said, every intermittent, even a quartan, may be readily cured. Two other formulæ are given in the same work*, and recommended as highly useful in the cure of intermittents, viz.

R. Tart. crud. ʒi. Arsen. cryst. ʒss. Pip. long. ʒss. Lap. prunell. ʒiss. Specif. febrifug. Crollii ʒiij. M.

The dose is from gr. v. to ʒss.

The other is

R. Arsen. alb. gr. v. Lap. prunell. vel Nitri depur. gr. xv. M. pro una dosi.

Professor Ackermann† relates, that in Pausa, a town of Saxony, a surgeon's family had been possessed for more than a century of a secret remedy against melancholy, which consisted of two grains of arsenic mixed with a drachm or more of white sugar, to be taken

* Dec. II. Ann. V. p. 474.

† Neues Magazin für Aerzte. Vol. II. p. 401. 8vo. Leipzig, 1780.

early in the morning, along with a large quantity of mucilaginous drink. The medicine produced a violent vomiting, so as to agitate the whole body, which continued not less than six hours; after this, he observes, the patient usually enjoyed a quiet sleep, and became more rational. The remedy was persisted in, care being taken that the effects of the first dose should be completely over before a second one was administered. Many repetitions of the medicine were not however requisite, as the disease, in general, soon yields to this mode of treatment; the patient was afterwards directed to continue a mucilaginous diet for a few weeks. Professor Ackermann examined some of the patients who had been cured by the surgeon at Pausa, and found that no ill effects had arisen in consequence of it. The same person, it is added, employed arsenic very frequently for the cure of intermittents; he dissolved two grains of arsenic in a pint of water, and gave two, or three table spoonfuls for a dose every day; under this treatment the fever seldom recurred more than twice; but he remarked that the patients were longer in recovering their strength than when the bark had been used.

Professor Ackermann farther observes, that another surgeon in the same place likewise

employed arsenic with great success; he gave fifteen drops of a solution of arsenic in water, along with alkaline salt, but the Professor had not been able to ascertain the exact proportions. A dose was ordered to be taken as soon as the patient felt the approach of the fit, and a quantity of warm tea was to be drank immediately afterwards. This produced a vomiting, which was encouraged as much as possible by repeated draughts of the tea. In this manner, it seems, he had cured many obstinate agues by two or three doses of the solution; and, amongst others, a quartan which had continued upwards of two years.

From some of the foregoing narratives, arsenic seems to have been used with as little precaution as emetic tartar; and since it appears, on good authority, not to have been productive of bad consequences, even in very large doses, we may be induced to lay aside that extreme anxiety with which we generally prescribe it; and may be encouraged to persist in the use of a remedy which, when prudently administered, is both safe and efficacious.

Many of our most active and approved medicines, as preparations of mercury and antimony, the squill, foxglove, &c., are capable
of

of producing as violent effects in the constitution, when given in too large a dose, as arsenic itself. All these medicines met with the same, if not stronger, opposition when first introduced, as arsenic does from many at present. It is well known that antimonial preparations were declared to be poisonous, and that the use of them was prohibited by a decree of the faculty of Physic at Paris in the year 1566; which decree was not repealed till 1637. We shall cease, however, to wonder at the prejudices formerly entertained against these medicines, when we consider, that even at the present day similar objections are made upon the Continent, especially in Germany, to the use of the bark, a remedy, the reputation of which has been so fully established by the united testimony of so many eminent practitioners, supported by almost innumerable experiments.

Mr. Theden, one of the most celebrated surgeons in Germany, and Surgeon General to the Prussian army, in speaking of the treatment of intermittents, observes*, that when his patients had previously enjoyed a healthy state of body, he was generally able to effect a cure in six or

* Unterricht für die Unterwundärzte. 8vo. Berlin, 1793.
eight

eight weeks. As he entertained the common idea that bark is apt to produce obstructions and enlargement of the viscera, œdematous swellings of the extremities, &c. he cautiously avoided giving this remedy until he had tried every other means. During the first three weeks he employed different medicines, with a view to loosen the morbid matter, and to render it fit for expulsion from the body; he then gave two ounces of bark, in doses of half a drachm, every two hours. After an interval of eight days, during which only bitters were prescribed, he ventured again to exhibit an ounce of the bark, and thus completed a cure. He cautions us against the use of bark whilst the face retains a yellow tinge, or whilst the febrile matter remains in the constitution; he confesses, at the same time, that he has seen œdematous swellings of the lower extremities after agues where no bark had been employed.

Dr. Vogel* is likewise of opinion that many cachectic diseases, particularly obstructions of the viscera, dropsy, jaundice, phthisis, tympanitis, coughs, asthma, hemicranium, deafness, cataract, vertigo, &c. are frequently the con-

* Handbuch der praktischen Arzneywissenschaft. 8vo. Stendal, 1781.

sequences of an improper treatment of intermittents; more especially when the cure has been attempted by astringents, arsenic, &c. or even by an unseasonable exhibition of the Peruvian bark, whilst the morbid matter still remains in the system.

The objections to the use of these medicines are so vague, that they appear to originate from popular prejudice and ill-grounded theories, rather than from any just practical deductions; they will therefore have little weight with those who are not contented with bare assertions, but make actual observation and experience the standards of truth.

Having frequently found the most beneficial effects from the mineral solution, and having never observed any ill consequences to arise from its use, I may presume to recommend a trial of it to surgeons practising in warm climates, and particularly upon the coast of Africa.

The high price of bark may sometimes prevent surgeons of ships from laying in, at their own expence, such a stock of this valuable medicine as will enable them to employ it freely in every case which requires its use. For notwithstanding the frequent complaints of several respectable surgeons in the navy, the quantity
of

of bark allowed by government to ships on foreign stations, is much too small; and most of the merchant ships trading to this coast are still more insufficiently provided.

Of the two most frequent diseases upon the coast of Africa, the remittent and intermittent fever, it is certain that the latter, though less rapid in its course, and apparently less dangerous than the former, yet for the most part occasions that irremediable injury to the constitution, which so often befalls Europeans trading upon this coast. There are few, even of those who are said to be seasoned to the climate by long residence, who have not suffered severely from repeated attacks of intermittents. This in a great measure arises from the unhealthy situation in which they live for the convenience of trade. They generally fix their residence on the banks of some river, or narrow creek, whose oozy shores, surrounded by mangroves, and excluded from the wholesome breezes, are a constant source of miasmata and contagion; to this must be added the debauched and irregular course of life which most of them lead. Though seasoned to the climate, as they suppose, their unhealthy fallow complexions and emaciated bodies, the frequent hectic flushings of the face,

face, swelled legs, &c. attended with obstructions and enlargement of the abdominal viscera, sufficiently indicate to every observer the flattered state of their constitutions. The ague probably still continues to return once a month or oftener, and harrasses them a few days, without being much noticed; for the severity of the disease seems to be considerably abated by its frequent recurrence, though its bad effects in the end are equally certain. As their appetite during the intermission is frequently keen, and even voracious, they flatter themselves that the constitution is not impaired by frequent returns of the disease; many also are negligent, from a confidence in the popular prejudice, that a cold fit shows the absence of danger.

In these cases, therefore, when the bark cannot be procured, or, as more frequently happens, when the patient has conceived a disgust for it, and cannot be prevailed upon to take it in a sufficient quantity, the mineral solution promises to be a safe and effectual substitute for it.

During the last rainy season I have had frequent

quent opportunities of exhibiting the mineral solution in intermittents with the same good effects as in the preceding year. Out of the number of cases which occurred in the present season, I have selected the two following, as being the only instances of quartans I have met with since I began to use the mineral solution.

CASE XX.

Sept. 11, 1793.—John Thompson, a mulatto, aged thirty years, was seized, about two months ago, with an ague, which returned every second day. After the second paroxysm he took an emetic, and soon after the operation of this, an opiate, which appeared to put a stop to the disease. A month ago he was again seized with cold shiverings, followed by an increase of heat, which terminated by a profuse sweat. The fit now returns every fourth day; the cold stage of which, commencing about noon, is very severe: the hot stage continues through the whole night, with violent head-ach, and towards morning is relieved by a profuse sweating.

ing. His appetite is pretty good; his body open.

Capiat Vesp. Antim. Tartar. gr. ij. cu. P. Ipecac. ℞j.
Cras incipiat fumere Sol. min. guttas xij. ter die.

20. The emetic operated well. He took the solution regularly for four days, and then omitted it, finding no return of his complaint.

30. He has had no return of the paroxysm, nor has taken any medicine since he left off the solution.

CASE XXI.

Sept. 8.—Anne Crankepoor, a black, aged twenty-eight years, has every fourth day, at noon, a severe cold fit of the ague, which continues near two hours, and is attended with violent rigors and pains of her bones; these symptoms are followed by a hot stage of long continuance, but which terminates by profuse sweating. She is affected, during the whole paroxysm, with violent pain of the head, stomach, and back, which also continue through the intermission, though with some abatement. She has taken an emetic and two anodyne draughts without any relief; and has

has had no stool for eight days. Her head-ach is at present very severe; her pulse quick and frequent; her skin hot and dry.

Capiat statim Camphor. gr. x. Tinct. Opii, guttas xxv;
Aq. font. ℥ss. et cras mane Sal. cathart. amar. ℥iss.
part. vicib.

9. She sweated profusely with the draught, and is much easier this morning. Her head-ach is considerably relieved; her pulse soft and regular. Both doses of the Sal. cath. amarus produced vomiting.

Capiat statim Ol. Ricini ℥i.—Repetatur Haustus li. s.

10. She could not yesterday retain the oil on her stomach, nor has yet had a stool. She passed an easy night, and feels no complaint this morning, excepting great languor and lassitude, with a sense of weight and fulness in the abdomen.

Capiat statim Calom. gr. v. Extr. Cathart. ℥j.

11. The pills operated gently three times; her bowels are much easier; she feels a slight pain of the head and general uneasiness, as if the fit was approaching.

Incipiat cras fumere Solut. min. guttas x. ter die.

13. The fit returned on the 11th at the usual time with great violence. The pain of her head and stomach was also very severe; she yet
feels

feels some pain of her stomach, with great restlessness and uneasiness. The solution has not been taken till this day.

17. The paroxysm returned at the usual time on the 14th, when she was affected with very severe head-ach and pains of the stomach and back, which still continue, being accompanied with great languor. She has taken only five doses of the solution since the 13th, and those not at regular times. She was very costive on the 15th, when she took

Calom. gr. v. c. Extr. Cathart. gr. xv.

which operated twice. She expects the paroxysm to-day.

Repetatur Solutio.

18. The paroxysm did not return yesterday, until six P. M.; the cold stage was very severe, and attended with great pain of the stomach and head; but these symptoms were much relieved by two grains of opium. She sweated profusely during the night, and feels a slight head-ach and pain of the stomach this morning, with languor and debility. Her body is open; her pulse natural.

Repetatur Solutio. Sumantur Opii gr. ij. urg. dolore Ventriculi.

20. She continues still weak and languid;

VOL. VI.

F

the

the pain of her stomach was wholly removed by the opiate.

Repetatur Solutio.

23. She has had no return of the paroxysm since the 17th, and makes no complaint but of debility; she is, however, able to walk about, and her appetite is somewhat better.

Omittatur Solut. min. Capiat Infus. Corticis Angust. ʒiij ter die.

Early in October she had entirely recovered her health and strength.

II. *An Account of the good Effects of a Solution of Sal Ammoniac, in Vinegar, employed, as a topical Application, in Cases of lacerated Wounds. By Mr. Henry Yates Carter, Surgeon at Kettleby, near Wellington, in Shropshire.*

IN the second volume of *Medical Facts and Observations**, I took occasion to mention, in a cursory manner, the good effects I had experienced, in lacerated wounds, from a solution of sal ammoniac in vegetable acid, em-

* P. 14.

ployed

ployed as a topical application; and which, in such cases, I observed, had seemed to promote the union of the parts and to moderate the discharge. As this mode of treatment is very different from that commonly in use, and I have had occasion to try it in many cases of bad compound fracture, and other lacerated wounds, in which there has been a tendency to sphacelus, I have been induced to make it the subject of a distinct paper, and for this purpose have selected the following cases, from a greater number, in which I have used it; and these, I hope, may be deemed sufficiently interesting to procure their insertion in a future volume of the valuable collection above referred to.

CASE I.

A poor man, named Ingram, aged upwards of eighty years, received an injury on his right foot, from a carriage passing over and lacerating it from the instep to the toes. The wound had been neglected for some days, when I was requested by a benevolent gentleman in the neighbourhood to visit him, and found the foot

sphacelated as high as the ankle, and the inflammation apparently extending still farther.

I began with scarifying different parts of the foot, by which means I gave vent to a considerable quantity of acrid ichor. The whole foot was then well covered with lint, continued to some distance above the disease, and directed to be kept constantly wet with a mixture composed of half an ounce of crude sal ammoniac dissolved in a pint of vinegar. Internally he took the bark in substance, liberally, with opium, as he had a disposition to diarrhœa.

On the second day after this mode of treatment had been adopted, I had the satisfaction to find that the inflammation had not sensibly increased, and that the patient felt at intervals a throbbing, but which, he said, was not painful, about the ankles. His pulse, which had been much quicker, was now at 100.

On the sixth day, a visible separation of the morbid parts was discoverable, and matter was perceptible on the verge of the separating parts; a fluctuation was felt in several parts of the foot, particularly beneath those places that had been scarified; and upon making deeper incisions here, we discovered a collection of good pus and granulations of new flesh. In the
course

course of a fortnight, the sloughs, having previously become loose, were gradually taken away, and the parts exposed one clear uniform wound. After this the bark was administered less frequently, but the use of the lotion was continued till the wound was nearly healed, which happened in about two months.

CASE II.

A girl, aged nineteen years, was attacked by a mastiff, and had the muscles of the thigh and leg, particularly the vastus externus and gastrocnemius so violently lacerated, that the worst consequences were to be expected from the circulation being cut off in the large vessels from the extremity, notwithstanding which she lost little or no blood; a circumstance, by the bye, that frequently occurs in lacerated wounds. She suffered but little pain, although the separated muscles of the upper part appeared to be much irritated. The large portions of muscle yet adhering were cautiously replaced as near their original situation as the nature of the case would admit; and after the parts had been well

bathed with warm vinegar, and due proportions of lint applied round the limb, the whole was encompassed with a broad roller, applied merely tight enough to retain the dressings; the limb was then laid in an horizontal position, and the pressure taken from the affected part by means of a pillow placed under the lower part of the leg, considerably below the injury. The whole was then wet with a lotion composed of half an ounce of crude sal ammoniac dissolved in a pint of vinegar, and ordered to be kept so constantly.

The first day she was but little sensible of the application. At night a draught, containing twenty drops of laudanum, was given, and she rested well.

On the second day I found her pulse but little quickened, and her thirst moderate; she had perfect feeling in every part of the limb, and complained of an acute smarting in the wound upon every renewal of the lotion, which continued for a few minutes, and then she became easy. An opening draught was given this morning, and she repeated the opiate at night.

On the third day matter seemed to be forming,

ing, but there was no appearance of inflammation or swelling of the limb.

On the fifth day from the receipt of the injury, the bandage was carefully removed, and I had the satisfaction to find that the muscles had united, and that the parts of the bone that had been laid bare were covered with new flesh. The discharge was kindly, and in moderate quantity, and the limb was free from pain. The same mode of dressing and the same applications were continued without alteration during three weeks, at the end of which time the cure was complete.

CASE III.

A young man, aged nineteen years, by a fall of coal in the pit while he was stooping, was pressed to the ground, and had his thigh broke about four fingers breadth above the patella. The upper part of the bone was forced through the muscles and into the ground, so that the hollow of the bone was filled with dirt, and stripped bare nearly four inches, and the muscles much lacerated. In this situation

he was brought home, (about a mile) and I then saw him ; the wound bled but little.

In this case I determined to try the effect of keeping the limb gently extended, nearly at its original length, after taking off so much of the bone as I should find requisite to a complete and exact reduction and to get above the coal flack which had been introduced.

As the bone was shivered longitudinally, I found it necessary to take off about three inches of it. This being done, and the wound well cleansed with warm vinegar and a small proportion of spirit of wine, I placed the lower part of the limb as exactly parallel to the other as possible, and retained it in that position by means of proper bolsters on each side of the limb. An eighteen-tailed bandage having been previously laid under the part, the dressing was made by gently filling the vacancy (the whole side of the leg *externally* being open) with soft pledgers of lint dipped in the same solution as that used in the preceding case, and the bandage was then applied as gently as possible, in order to prevent the flesh from being pressed into the part that the bone ought to have occupied ; and a splint applied externally on each side, merely to give more steadiness to the

I

limb,

limb, but without occasioning much pressure. I think it right to mention also that the middle tails of the bandage were cut smaller than the others, and applied in such a manner that the wound might be uncovered, in order that the lotion might be applied immediately to the wound, without disturbing any other part.

He was let blood, and twenty-five drops of tincture of opium were given at night, and the attendant was strictly enjoined to keep the part constantly wet with the solution, except only during the intervals of sleep.

Upon visiting him the morning after the accident, I found he had had but little sleep, though his limb had given him but little pain, except for about a quarter of an hour after the application of the lotion, after which he said he had felt the whole leg and foot become sensibly warmer. The lower part of the limb lay very steady, exactly in the situation in which it had been placed; he took this morning three grains of calomel, which procured one stool.

On the fifth day, including the day of the receipt of the injury, (there having been some appearance of matter between the folds of the bandage) the dressings were wholly removed, and the wound was found covered with a well-concocted

concocted pus in moderate quantity, and with new granulations. The dressings were continued in the same manner as before, the whole vacancy being carefully filled with dossils of lint, made as soft as possible, till the whole was level with the skin; and over these the bandage was applied as before. He continued to repeat the opiate every night, and the calomel occasionally; his appetite was tolerably good, he used nearly the same diet as when in health, and was permitted to drink a small quantity of ale.

On the eighth day the dressings were again removed, and the appearances continued to be favourable. From this time, the weather being warm, the wound was dressed every day in the same manner as at first; and in about eight weeks the callus was completely formed, and had filled up the void space, and the wound was reduced to about a quarter of an inch in diameter.

In ten weeks he came down stairs, and went about on crutches; and in about sixteen weeks from the time he received the injury, he went with a stick only, and was able to walk nearly two miles. The limb was not quite an inch shorter than the other; the small ulcer continued

nued to discharge, till a considerable exfoliation of bone, which gradually made its way outwards, was extracted, after which the wound soon healed.

CASE IV.

A boy, aged about fifteen years, had the misfortune to slip his hand under the axletree of a water-wheel, which moves at about the distance of two inches and a half from a brick wall or buttress supporting another building; his arm was taken in to the elbow, and the machine performed several revolutions on the part before he could be extricated. The flesh was stripped down on each side of the thick part of the arm, and the thumb was nearly separated; but the fingers and hand had suffered but little, and there was no hæmorrhage. The thumb was not taken off, but carefully replaced, as well as the other muscular parts that had been separated; and to the whole wound a large quantity of lint was applied, wet with the solution of sal-ammoniac in vinegar. He took twenty drops of tincture of opium at night,
but

but he was very restless, and complained much of his arm.

Second day. The arm had bled in the night, and the dressings were become stiff and hard, which rendered it necessary to remove them. The disturbance this occasioned produced a degree of inflammation which, I believe, might otherwise have been prevented, and which proved the source of misfortune. The parts from this time became excessively painful, and the inflammation extended to the upper part of the arm, and to the shoulder and side, as far down as the pectoral muscle. He was costive and feverish, and complained much of thirst. The whole arm was wrapped in a cataplasm made of oatmeal, with equal parts of vinegar and water; and three grains of calomel were immediately given. Two stools were procured by this medicine; but the pains still continued to be very distressing to him. His pulse was at 100.

Third day. The above symptoms continued; the pulse was increased to 110; and he was at times delirious; the upper parts of the arm, shoulder, and side, were become of a dark red colour, and were exceedingly tense. He had several loose stools; the arms and side were
dressed

dress'd as before, with the addition, in the liquid of which the poultice was made, of half an ounce of crude sal ammoniac, and an ounce of spirit of turpentine. He took half a drachm of Peruvian bark, with fifteen drops of tincture of opium, every third hour; and care was taken to distil some of the solution between the dressings, upon the shoulder, very often, in such a manner that it might make its way to the affected parts.

Fourth day. I found the whole fore arm, from the elbow, completely sphacelated and dry; but the shoulder and side were nearly in the same state as yesterday, the inflammation not having increased; his purging had ceased; he was not so thirsty, and his pulse was at 100; but he complained much of head-ach and weariness. Notwithstanding there appeared some reason to conclude that his head-ach might, in some measure, be occasioned by the quantity of opium he had taken, I continued the use of it in the same doses; a stool was procured by means of a clyster. The use of the lotion was continued.

Fifth day. The symptoms were nearly the same as yesterday. The same dressings and medicines were continued.

Sixth

Sixth day. The pain and tension were much lessened, he had rested tolerably well, and was free from thirst; the shoulder and side, with a considerable part of the upper arm, seemed approaching to their natural colour, and the extent of inflammation was visibly decreasing. The bark was still continued, but without the tincture of opium, instead of which he took two grains of purified opium at night.

The cataplasim was continued as before for about a week, from this time, when the shoulder and side having recovered their original tone, it was changed for one composed of oatmeal and the solution alone. In a few days matter formed plentifully round the bone in those parts where the lacerations had been deep, and large portions of the muscles were cautiously removed. The matter formed was of a good consistence, and moderate in quantity; and the wound was perfectly easy, excepting only upon the application of the lotion, and for some short time after. The whole hand dropped off at the wrist; the other parts gradually filled up with good flesh, and are now completely healed.

CASE

CASE V.

A man, aged thirty-six years, by the fall of a very heavy iron rod perpendicularly upon his foot, upon that part where the shoe is generally buckled, received a considerable lacerated wound, by which the tendons were much injured, and the integuments and muscular flesh were stripped off from the upper part of the tarsus, and hung in a large loose flap down the side of the foot. The wound bled considerably, and the whole foot, from the violence of the blow, was insensible. The parts were well cleansed from the grumous blood with vinegar and water, with a small quantity of spirit of wine, and the loose flap replaced in the situation from which it had been torn, and dressed with pledgets of lint dipped in the solution; and a cataplasm applied of oatmeal and vinegar.

The morning after the injury, upon removing the dressings, the wound and whole foot were found to have a favourable appearance; but at night he began to complain of a great degree of heat, throbbing, and sense of tension.

On

On the third day, on removing the dressings, the whole upper part of the foot appeared to be hastily approaching to a sphacelated state. It had lost all sensibility to the touch, and the inflammation had increased, though in so short a time, considerably above the ankle, and to the extremity of the toes. A sensation of burning heat in the whole foot and leg still continued. The parts that were loose were now removed, and the wound, after having been bathed a considerable time with a mixture of warm vinegar and water, with a small quantity of sal ammoniac previously dissolved in it, was dressed as usual, the lint being first well saturated with the lotion; and over the whole a cataplasm was applied as before. A purgative medicine, composed of four grains of calomel, and five grains of aloes, was given, which operated well. He passed this day with somewhat more ease, and at night took thirty drops of tincture of opium.

Fourth day. He complained of having passed a very restless night, and that the painful sensation of burning heat still continued; the inflammation went on increasing; his pulse was at 97, and he had much thirst and flushing eat. Bark, in the quantity of half a drachm,
was

was given every third hour, and twenty drops of tincture of opium every sixth hour. The same dressings were continued, with the poultice; but at night the poultice was omitted, and the dressings kept wet with the solution alone.

Fifth day. He had rested much better; his thirst was more tolerable, and the heat and other symptoms were much more moderate; his pulse was at 90; the inflammation had not increased; and the tension about the ankle was lessened. The same medicines and local applications were continued as last night. On renewing the dressings in the evening, he complained of having passed a very painful afternoon, and that the sense of heat had been greater. He attributed all this to the omission of the poultice, which was now, at his earnest request, renewed.

Sixth day. In the morning the symptoms were much increased, and the inflammation was spreading, with a violent degree of pain and tension, the whole upper part of the foot being in a sphacelated state; and the patient complained of excessive pain. The same dressings as before were applied, but without the poultice, after bathing the parts with warm vinegar; a broad roller, for the convenience of

keeping the parts wet, was gently applied over all the inflamed parts; and as I had a suspicion that the increase of his pain, &c. yesterday, if not wholly, was, in a great measure, owing rather to a want of due care in keeping the parts constantly moist, and thus suffering them to get dry and hard, than to any effect the application could have in producing those symptoms, I paid this day a particular attention to this circumstance, by visiting him several times, to see that the solution was duly applied; and in a few hours the symptoms of pain and heat in the whole limb were greatly diminished, and continued gradually to abate the whole day. His pulse at night was at 93.

Seventh day. The symptoms were nearly the same as yesterday; the inflammation, upon the whole, was rather less, but there was no appearance of matter. He had passed a tolerable night; but his pulse was still at 93. As he was costive, the purgative medicine was repeated.

Eighth day. He had past a good night, comparatively speaking; the pain in the upper part of the limb (or above the disease) was considerably lessened, and the inflammation was much less; a small quantity of matter appeared upon the edges of the lacerated parts; his

his pulse was at 90. He began to complain of severe smarting upon the renewal of the lotion, and at times insisted on its application being deferred to longer intervals, though when the parts began to grow dry, the heat and sense of stricture were constantly renewed.

Ninth day. He had passed a restless and painful night; his foot and leg were in much pain at intervals, but (exclusive of the smarting pain for a quarter of an hour upon the lotion being applied) he always became much easier after the wetting of the parts, which took place once in about two hours, unless sleep intervened.

From this time the use of the lotion was continued in the same manner as before, and he continued also to persevere in the use of the bark and opium; the sloughs separated kindly; the inflammation went off from the leg and toes, and a separation of the diseased parts took place at a very little distance from the edges of the original injury. The wound discharged a well-formed matter, and as the parts beneath some of the thickest sloughs granulated, the latter gradually came away without much pain, and the whole was healed in ten weeks, except a very small ulcer upon the lower part

of the Tarsus, through which a small exfoliation made its way.

As in the preceding cases I was careful to obviate the effects of irritation, by keeping the bowels moderately open, giving occasionally, and sometimes liberally, of opium; and invigorating the system by means of wine and the Peruvian bark; it may perhaps be suggested, by some readers, that the favourable termination of the cases I have been relating was due rather to the internal than external remedies employed; and that to subject to a fair and decisive trial this or any other remedy, no other should be employed at the same time. This is indeed what I have done in slighter cases of laceration, where local applications only were requisite; and in all such cases the union of the parts has appeared to me to be much more speedily effected by means of the lotion, than it is by the ordinary mode of treatment. And I am able to recollect no instance of bad compound fracture, or of lacerated wounds, attended with or threatening sphacelus, where the warm fomentations and cataplasms commonly employed in such cases were

were made use of, in which there was any such obviously good effect from the local treatment, as in the cases I have been describing; notwithstanding there was the same liberal use of opium and Peruvian bark, &c. internally. On the contrary, I have but too often seen the worst effects from such cataplasms, &c.; and in one of the above cases, (Case V.), the bad effects of a poultice, applied at the earnest request of the patient, were very striking, when contrasted with the relief he afterwards experienced from the use of the lotion.

III. *Case of a diseased Kidney. By the same.*

A SEAMAN, forty years old, of a plethoric habit, applied to me at Port Royal, in Jamaica, in 1782, with complaints nearly as follow :

A constant aching, and sometimes acute pain, about the region of the right kidney, attended with a numbness of that side, and pricking pains

pains along the urethra, particularly when he passed his urine; frequent inclination to make water, sometimes without ability to void any, and never voiding it but in small quantity; the urine itself being high coloured, depositing a gritty lateritious sediment, smelling very strong, and forming a film on its surface, which approached to a yellow colour. He complained likewise of a sense of fulness and heat at the neck of the bladder and about the perinæum, and could get but little rest in any other than an horizontal posture. He was costive, and had frequent nausea.

As he had a full pulse, ten ounces of blood were taken from the arm, and a purging draught was administered; after which he took occasional doses of a mixture, the principal ingredients of which were diuretic salt and tincture of opium.

In the course of two or three days his pain was much alleviated, but the difficulty with which he voided his urine still continued.

He now complained of frequent and painful erections, more especially when an inclination to make water came on; he had likewise profuse colliquative sweats, and was costive.

Care was taken to obviate this disposition to
costive-

costiveness, by means of purgative medicines and clysters. Opium was now more liberally administered, and recourse was occasionally had to the warm bath. This last produced a certain degree of ease while he remained in it, but the sense of stricture about the neck of the bladder continued, and the quantity of urine he was able to void seemed every day to become less, so that at the end of a fortnight it was deemed necessary to make use of the catheter, as he was unable to pass a single drop of urine without it.

By means of this instrument, from four to six ounces of turbid urine were drawn off twice a day. He had now much fever, and the pain about the neck of the bladder was become very acute, and seemed to affect him spasmodically, as well after as previously to the introduction of the catheter. He was likewise frequently seized with violent pain, which began in his shoulders, and proceeded along the right side to the hip.

About a month after the first use of the catheter, he complained of a pain in the urethra, near the seat of the prostate gland, particularly when the instrument was passing; and

at times the catheter seemed to meet with some resistance at that part.

From this circumstance, together with the continuance of the pain in that and the neighbouring parts, and the frequent discharge of drops of a mucous consistence from the urethra, we were inclined to think that the principal seat of the disease was in the prostate gland, (especially as no appearance of calculus had been observed), when a fresh set of symptoms directed our attention more particularly to the right kidney.

These symptoms consisted in a pain about the region of that kidney which he had before scarcely mentioned, but which now (about seven weeks after he first made his complaints known) was, at times, very severe. His shoulders also, but particularly the right, were sore, and at intervals acutely painful; the inguinal and axillary glands became swelled, and sore to the touch; and he complained frequently of a sense of coldness in the direction of the right ureter, which was succeeded by a painful inclination to make water.

From these circumstances it was suspected that the right kidney, if not the chief source of the extraordinary symptoms I have been describing,

describing, had at least suffered considerably. He was therefore urged to recollect any external injury he might have received. After a little hesitation he informed us, that about a month previously to his first applying for relief, he had received several violent blows from the end of a large rope across his loins, which for some time had given him considerable uneasiness. In the course of a few days, however, he said, the pain had gone off, but had returned at intervals; and as he had suffered much, at different times, from gravel, he had ascribed his present complaints to that cause.

At the time he made known these particulars, he was in a very reduced condition; his stomach was become so extremely irritable, that it retained but little of what was given to him either of food or medicine; and about a week afterwards he died.

On dissection the urethra was found to be in a healthy state, but the prostate gland was a little enlarged. The bladder contained about eight ounces of turbid urine, mixed with a purulent fluid, very offensive to the smell. The right ureter was much enlarged, and filled with the same kind of foetid matter. The kidney

ney on the same side was enlarged nearly to thrice its natural size, and on being opened was found to be in a state of suppuration, and to contain a considerable quantity of foetid pus, so that the internal substance of the kidney was in a great measure destroyed.

There was no appearance of calculus; and the other kidney, as well as the rest of the abdominal viscera, appeared to be in a natural state.

It may be doubted, perhaps, whether the affection of the kidney, in this case, ought solely to be attributed to the effects of the blows that were inflicted; but allowing the kidney to have been previously diseased (and the complaints the patient had already experienced, and which he attributed to gravel, render it not improbable that it was so); still there can, I think, be no doubt that the suppurative process which took place was hastened, if not immediately occasioned, by external violence. And of suppuration of the kidneys from external injury, in any respect similar to the present, I have been able to meet with no example in books. Different systematic writers do indeed
 enume-

enumerate external contusion among the remote causes of nephritis, but I do not find, in any of them, an instance of such an affection from such a source; so that I flatter myself the case I have related will be thought worthy of being recorded.

It shows that a frequent inclination, without ability, to make water, is not always occasioned by gravel or calculous concretions; and it affords a striking instance of the influence an organ like the kidney may have upon parts not only contiguous to, but even remote from the seat of disease.

IV. *Case of a Gun-Shot Wound of the Head. By the same.*

A HESSIAN grenadier, aged between thirty and forty years, being one of a detachment sent to reduce a fort on the banks of
the


the Delawar, in the act of levelling his piece, received a ball (grape shot) on that part of the os frontis which forms the external canthus of the eye. The ball making its passage through the head, came out under and rather behind the opposite ear, as in the annexed plate*.

What were the immediate effects upon the receipt of the injury I am not able to say, not being immediately upon the spot; but he appeared, when brought to the regimental hospital, to have a perfect recollection of every circumstance that had occurred to him, except only for a short time after he fell. He complained of little pain, and did not appear to have lost so much blood as might have been expected.

The ball being a spent one, had much splintered the cranium, both at its entrance and exit; and was found in the folds of his coat collar.

The wounds being cleansed, and the splinters of bone removed, as far as was practicable, from about the external parts, suitable

* See Plate I. Fig. 1. in which *a* refers to the entrance of the ball, and *b* to the part where it passed out.



dressings were applied; and his pulse being full, he was let blood; after which he took twenty-five drops of tincture of opium. The next day he had a sense of heaviness over his eyes, and observed that objects did not appear to him so brilliant as usual; towards the evening he complained of nausea and thirst. He took *tart. vitriol.* and *antim. diaph.* $\bar{a}\bar{a}$ gr. xii every third hour, and a clyster was administered. On the third day he complained of pain of his head, accompanied with drowsiness; and, at intervals, of a weakness of his extremities. As the clysters had failed to procure a sufficient discharge of fæces, he was directed to take three grains of calomel and fifteen grains of powder of jalap, which operated well, and procured an alleviation of the symptoms just now mentioned. His eyes were but slightly inflamed, and he complained of but little pain in that on the affected side.

On the 6th day there was a good discharge of matter from the wound, and escars began to separate in pretty large sloughs. From this time he rested tolerably well without the use of the opiate, which till now had been repeated at bedtime. Splinters of bone, that had been driven in at the superior wound by the ball, came
away

away from the dependent orifice at almost every dressing (which was twice a day) for several days. The nausea, head-ach, weakness of his limbs, thirst, and every symptom of fever, gradually vanished; the superior orifice filled up with new granulations, and cicatrized firmly; and in about ten weeks there remained nothing more necessary than a superficial dressing to the inferior opening near the ear.

I did not see this man after he had actually left off every application to the affected part; but from the condition of the wound, and the patient's health and vigour, I have not any room to doubt, that in a few days, after I last saw him, he was capable of returning to his duty.

On reflecting on this extraordinary injury, (inasmuch as it was not a mortal one) I am inclined to think, that as the ball, though a large one, entered low down upon the orbit, and near the external part of the eye, it missed the os planum and frontal sinuses, and consequently that branch of nerves that passes through them; so that, judging from its apparent direction, it must have injured part of the os ethmoides, near the septum nasi. To this course of the ball,

ball, and the favourable situation of the dependent orifice, the favourable event of the case was probably owing; for though he complained at certain periods of a sense of weight upon the upper and fore part of the head, general weakness of his limbs, and loss of sight, symptoms indicating an oppression of the brain, yet upon opening the wound, and giving vent to the matter, which was in some measure confined by the dressings, those symptoms gradually vanished, and the patient always became perfectly easy after the application, for a few minutes, of a warm fomentation.

An instance of a ball entering under the right eye, and passing obliquely through the cerebrum and cranium above the right ear, without hurting the eye or sight, is recorded by Heister in his *Medical, Chirurgical, and Anatomical Cases and Observations*, page 7 (of the English translation) Obs. VII.

V. *An Account of some extraordinary Symptoms which were apparently connected with certain morbid Alterations about the Veins and Nerves. Communicated in a Letter to Dr. Simmons by Mr. John Pearson, Surgeon of the Lock Hospital, and of the Public Dispensary.*

MRS. P. aged fifty-one years, of Miles' Lane, Cannon Street, began to suffer from a peculiar uneasiness at the inner part of her left leg, about seventeen years ago, when she was in the third month of her second pregnancy. The skin which covered the particular seat of her complaint, retained its natural colour; but there was a circular induration, of about half an inch in diameter, very little elevated above the surface, which was exquisitely painful when slightly touched or compressed; this morbid part was situated in the course of the vena saphena major, and was about six inches above the joint of the ankle. Besides the acute pain which was produced by inadvertently touching this little tumour, Mrs. P. commonly suffered several paroxysms of
of

of pain every day; each of these attacks was accompanied with an increased redness, and a sensible elevation of the indurated part, the pain at the same time extending to the knee, and often darting to the stomach; the duration of the fit was about twenty minutes; it was attended with slight convulsive motions of different parts of the body, and frequently terminated with flatulent eructations. These fits of pain did not recur at any regular periods; so that the number which she underwent in the course of a day was various and uncertain; for a disordered state of the stomach, or a sudden perturbation of mind would at any time excite one of the paroxysms. She also had observed, that the severity of her sufferings was invariably increased during the periods of menstruation and of pregnancy; and that in the latter months of gestation, the duration of each recurrence of pain was extended to an hour and a half. But although this disease was uniformly aggravated by certain alterations in the state of the uterus, yet it continued with undiminished severity after Mrs. P. had ceased to bear children; for when her youngest child was more than six years old, she had not experienced any abatement of her daily sufferings. About thir-

teen years ago, I advised her to have the morbid part removed; but at that time she was unwilling to undergo an operation; she however submitted to various methods of treatment, under the direction of different medical gentlemen, but without obtaining any relief.

In the month of April, 1793, Dr. Lowder, who had been long acquainted with the circumstances of this painful complaint, informed Mrs. P. of the success which had attended the removal of a similar tumour, by the application of a caustic. She read the history of the case, which is published in the third volume of the *Memoirs of the Medical Society of London*, and very soon determined to seek relief from the same mode of treatment.

Accordingly, on the 22d of April, I applied the lapis infernalis to the morbid part; she endured the most excruciating tortures during several minutes after its application; but the pain gradually diminished with the sensibility of the part, so that in about twenty minutes the eschar was completely formed, and she then felt no more pain than what is the usual consequence of a caustic applied to any part of the body. From this day she never experienced the recurrence of a single paroxysm of pain; the eschar

exfoliated in about twelve days; and on the 7th of June the fore was perfectly healed.

As the preceding history contains some curious and rather uncommon circumstances, I beg leave to offer a few observations upon some of them. The indurated part having been destroyed by a caustic, it was not in my power to examine its internal structure, so as to discover the true nature of the morbid alteration. I ascertained, however, that a portion of the vena saphena major, and that branch of the crural nerve which accompanies it in its course down the inside of the leg, were completely included within this tumour. This fact was clearly demonstrated after the exfoliation of the eschar; for I then saw a portion of the vein hanging down at the superior part of the fore, and the naked nerve in contact with it; and on touching the nerve with my probe, Mrs. P. instantly complained of an acutely painful sensation, similar to that which she had been accustomed to feel before the tumour was removed. I then destroyed that part of the nerve which was exposed with lunar caustic, and my patient suffered no more uneasiness. After thus proving

that a vein, and a considerable ramification of a nerve, were contained within the diseased part, I proceed to observe, that the paroxysms of pain were excited by every thing that accelerated or otherwise disturbed the circulation of the blood; whether applied to the induration, or affecting the general system; as all strong exertions of the muscles, external impulse, or mental commotion. The ascent of the blood, in the veins of the lower extremities, is necessarily impeded in the state of pregnancy; and during this period, the fits of pain were always sharper, and were also of longer duration; and at the time of parturition, when the action of the heart and blood-vessels is considerably increased, Mrs. P. suffered exceedingly; for, to use her own expression, she “had all her labour pains in her leg.”

It is also highly probable, that the portion of vein which passed through the tumour was unusually distended with blood at the time of the paroxysm; for upon these occasions, the morbid surface became redder than common; and the tumour was sensibly elevated. We may therefore, perhaps, venture to conclude, that the vein and the nerve being confined within a substance that could not be easily distended, when-

ever

ever the vein became preternaturally turgid, the nerve was compressed between its parietes and the internal surface of the induration; and th at consequently the symptoms were connected wi th this state of the part. I do not suppose that it will be necessary for me to undertake a proof in detail, that a certain degree of pressure upon a nerve will produce pain, spasms, and convulsions; it may be sufficient for my purpose to refer to a few of the many instances which are recorded in medical books. In the fourth volume of the Edinburgh Medical Essays, Dr. Short has related the history of an epilepsy, which was caused by the pressure of a hard cartilaginous substance upon a nerve; he cured his patient by removing the tumour, and dividing the nerve. Guattani, in his *Treatise de externis Aneurysmatibus*, (Hist. XX.) has recorded a case in which violent spasms were occasioned by the pressure of an aneurism upon a nerve. In the *Essays and Observations Physical and Literary*, Vol. III., the late Sir John Pringle has published a Case, where a tumour formed by extravasated blood, by pressing upon the intercostal nerves, produced pain, irritation, and perhaps a hic-

cup, which could not be stopped*. I do not intend to deduce any general conclusion from a particular instance; for although the remarkable symptoms which occurred in Mrs. P.'s case, were connected with a morbid state of a vein and a nerve; yet as no account has been published of the internal structure of parts which have been affected by a similar complaint, it would be improper to conclude, that every instance of local morbid sensibility, accompanied with convulsive motions and pain, must depend upon such a peculiar condition of the suffering parts. I have indeed seen another case, very much resembling that of Mrs. P.'s, in which there is a small exquisitely sensible induration

* For instances of convulsive motions, and even epilepsy, produced by local diseases about some of the extremities, or that were cured by the removal of matter, carious bone, or some portion of the integuments, consult Willis *de Morbis Convuls.*; Riverius *de Epilepsia*; Schenckii *Observat.* (*Lib. de Epilepsia.*); Forestus *de Cerebri Morbis*, Lib. X. Obs. 67; Petri Borelli *Hist. & Observat. medico physicarum*, Cent. II. Obs. 95. Joh. Rhodii *Observ. Med.* Cent. I.; Tulpii *Observ. Med.* Lib. IV. Cap. 2; Boneti *Septuagintarum*, Lib. I. Sect. 13; Van Swieten *Comment. in Aph. H. Boerhaave*, Tom. III. § 1075. Haller *Elementa Physiologicæ*, Tom. IV. § 30. Simson on the Vital and Animal Actions, Essay I. ch. 3.

at

at the posterior part of the leg, near the beginning of the tendo achillis, from which the patient suffers acutely whenever it is touched. She has occasional paroxysms of pain, but they return at uncertain intervals; and she thinks that they grow milder. In this instance, as in that recorded by Dr. Bisset *, the tumour becomes uneasy in rainy and windy weather; but it does not appear that the disease had ever any connexion with pregnancy. I suspect that the tumour, which I have just now mentioned, may be connected with the vena saphena minor, and that consequently it may include or compress a small branch of the sciatic nerve; but as I could not render the cutaneous veins of the leg turgid by moderate pressure, its exact situation was not ascertained †.

In

* Memoirs of the Medical Society of London, Vol. III. Art. VI.

† The first volume of M. Pouteau's posthumous works contains a very curious history of a disease which he there calls cancerous; whether properly or no I shall not inquire; but as it resembles Mrs. P.'s case in some of its characters, I shall take the liberty of presenting an abstract of it:

“ On voyoit à la partie basse du *Sternum* une surface
“ ovale de largeur d'un ecu de six livres dans son petit dia-

H 4

“ metre,

In the early part of the last Spring, a young married woman applied to me at the Public Dispensary, complaining of pain and lameness of the right arm. She shewed me a tumour of a pale red colour, and of about the size of a filberd,

“ metre, sans élévation, sans rougeur, sans engorgement
 “ circonvoisin. La peau seulement qui la recouvroit étoit
 “ un peu moins nette, que par tout ailleurs, mais semblable
 “ à la sensitive qui paroît craindre la main qui l’approche.
 “ Cette portion des tégumens auroit fait ressentir les plus
 “ vives douleurs, si le doigt, sans la toucher, en eût ap-
 “ proché avec trop de célérité. Le moindre insecte, un
 “ fœtu que le hasard auroit fait poser dessus, eussent aussi-
 “ tôt rappelés les convulsions. Les rétors de ces convul-
 “ sions étoient périodiques, se montrant à sept heures &
 “ demie précises du soir. Dans le plus grand calme, on ne
 “ les attendoit que de deux jours l’un; & à la moindre agita-
 “ tion, les mouvemens convulsifs étoient journaliers. Leur
 “ durée étoit de deux heures, & même plus.” The his-
 tory presents us with many other extraordinary circum-
 stances; but it may be sufficient at this time to add, that
 M. Pouteau made a crucial incision in this morbidly sensi-
 ble part, which afforded an immediate although but a tem-
 porary suspension of the pain and convulsions. He then
 extirpated the portion of diseased integuments; but as the
 young lady was not perfectly relieved by this operation, he
 finally completed the cure by burning a cylinder of cotton
 upon the part. Vide *Oeuvres posthumes de M. Pouteau*,
 Tom. I. ch. 1.

which

which was situated in the course of the vena mediana basilica, at the bend of the arm: this morbid part was constantly uneasy; but when it was pressed or handled, she complained of acute pain, which extended along the upper arm, and produced slight convulsive motions in the muscles. She derived no advantage from mild discutient and emollient applications; but her pain increased so much, that her health became injured, and she was at length confined to her bed. On visiting her at home, I found the tumour unaltered in its appearance, excepting a spontaneous separation of the cuticle from its surface; she was in constant pain; the uneasiness not only proceeding along the upper arm, but also to the neck, and affecting the breast and muscles on the right side. Her pulse was feeble, but not too frequent; she complained of a great sense of weakness, and convulsive motions were excited in the muscles of the upper arm, neck, and thorax, on that side, by the gentlest examination of the morbid part. I ordered a large vesicatory to be applied on the inner part of the fore arm, and directed her to take ten grains of *pulvis ipecacuanhæ compositus*, whenever her pain should be unusually severe. She soon derived considerable relief from this mode
of

of treatment : the blistering plaster was repeated twice during my attendance ; the tumour gradually became less painful, and diminished in bulk ; and in about a month it had entirely disappeared. It was not more than three weeks after she was dismissed, when she applied to me again, on account of a tumour very much resembling the former one, which was situated at the bend of the arm, in the course of the *vena cephalica* ; so that a portion of the vein evidently passed through, or, rather, was included within the center of the morbid part. The pain and morbid irritability affected the same parts as before, but in a much inferior degree. I directed a mode of treatment similar to that which had been employed on the former occasion, and it was attended with equal success.

This young woman had some symptoms which indicated a diseased state of the lungs ; and she occasionally spat blood : but she had not been formerly subject to any particular complaints ; she menstruated regularly ; and had never been pregnant. I cannot assign any probable cause for the appearance of so singular a complaint as that which I have now described ; but some of the effects which took place would
perhaps

perhaps admit of an explanation, if it could be proved that a small ramification of a nerve, as well as a portion of a vein, were included within each of the tumours. That this was actually the case is highly probable, because the cutaneous nerve distributes several of its branches in the vicinity of the vena mediana basilica; and small fibrils belonging to the musculo-cutaneous nerve, are commonly seen near the vena cephalica, and the vena mediana cephalica; so that tumours situated at the bend of the arm, and in the course of these blood vessels, must be almost necessarily in contact with one or more branches belonging to the internal, or external cutaneous nerves*.

* The late Professor Camper, in a valuable work entitled *Demonstrationum Anatomico-Pathologicarum Liber primus continens Brachii humani Fabricam et Morbos*, has given a very distinct view of the mode in which these small branches of nerves are distributed at the bend of the arm; and his engravings are accompanied with some good practical observations. Mr. Abernethy also published two engravings, last year, in the second part of his *Surgical and Physiological Essays*, in which the course of these nerves is very neatly and correctly delineated: and the essay to which they are annexed, contains many useful remarks “on the ill consequences sometimes succeeding to venæ-section.”

I beg

I beg leave to refer it to the intelligent reader, how far the following account of a disease of the subcutaneous nerves, as described by Professor Camper in the work already referred to, bears any resemblance to the preceding histories.

“ Non raro in nervis cutaneis tubercula par-
 “ va ac dura observantur, quæ vera ganglia
 “ sunt, pisi magnitudinem licet non excedant ;
 “ dies tamen noctesque acutissimis lancinanti-
 “ bus doloribus ægros torquent : externis re-
 “ mediis non cedunt ; scalpello igitur ea attin-
 “ gere oportet. Franequeræ ex cubito feminæ
 “ tale, plagâ factâ, sustuli, quod ramo musculo-
 “ cutanei nervi adhærebat : post operationem
 “ optime se habuit. In subcutaneis nervis fre-
 “ quenter esse videntur. Amstelædami simile
 “ ganglium genu mulieris occupans, eodem
 “ modo sanari curavi. In viris plus semel ea
 “ vidi : albicant intus, cartilagineæ duritiæ
 “ sunt, renitentia, & intra nervorum tunicas
 “ sedem habent.” Lib. I. P. 11. Cap. 2. § 5.

I have seen many symptoms resembling those which occurred in the preceding cases, apparently follow, as consequences of wounds inflicted on small branches of nerves ; but as this paper is already much longer than I expected it would have been, I must defer giving an account

count of them to another opportunity. As the following case exhibits some uncommon circumstances, I insert it as a kind of supplement to the foregoing histories.

“ The singular effects of an issue in the inside of the thigh, which appeared in the case of a clergyman ; written by himself, August 25th, 1793.

“ The Rev. Dr. T——, of Knightsbridge, above 60 years of age, having had a hint from a medical friend, that an issue might be of use to his health, he had one made by a blister, in the lower part and at the inside of his right thigh, about the end of May last. Two days after the pea was put in, he was seized with a sickness and vomiting, which continued several hours. In about six days after this first attack, he had a return of the same symptoms ; and these fits recurred every six or seven days. But what is very remarkable, when the issue began to discharge, he became deaf in both his ears, and the deafness arrived to such a degree, that in preaching he could but just hear his own voice.

“ After the issue had been kept open six weeks, it occurred to him, that perhaps the regular fits of sickness and vomiting, and the
“ unusual

“ unusual deafness, (both of which he recol-
 “ lected had commenced with the issue) were
 “ occasioned by a sympathy of the nerves ; and
 “ having made observations for one week longer,
 “ which confirmed this opinion, he determined
 “ to dry it up. This he did gradually, by using
 “ pease of a smaller size, till the ulcer was not
 “ more than one eighth of an inch in diameter.
 “ When the pea had been out only twelve hours,
 “ he was sensible of some small return of his
 “ hearing, and on looking at the sore, he found
 “ it healed ; which he considered as a farther
 “ confirmation of his opinion, respecting the
 “ cause of his deafness, as well as of the sickness
 “ and vomiting. He found, that as the wound
 “ healed, the deafness lessened, and when it was
 “ completely healed, his hearing was quite reco-
 “ vered, nor has he had one fit of sickness since.”

When Dr. T—— related his case to me, I
 desired him to let me see the cicatrix of the
 issue ; and on carefully examining it, it ap-
 peared probable that the pea had pressed against
 the side of the vena saphena. I would also far-
 ther add, that my examination of the part ex-
 cited a slight degree of nausea.

VI. *An Account of the Extraction of an extraneous Substance from the Rectum. By Mr. William Blair, Surgeon of the Lock Hospital; and of the General Dispensary in Newman Street, St. Mary-le-bone.*

ON Tuesday, the 25th of March last, a French gentleman was sent to me by an Apothecary in this neighbourhood, complaining of a pungent, hot, and irritating sensation in the *rectum*; which was considerably augmented during every evacuation *per anum*. These painful symptoms had commenced on the preceding Sunday, and continued to encrease in so alarming a manner, that, upon the day following, he was induced to examine with his finger, whether or not any foreign substance, or other cause of his uneasiness, could be discovered in the intestine. He had the good fortune to feel something in the rectum, which he thought was unnatural, but could not remove it; and therefore he applied the next day for surgical assistance.

Having submitted the patient to a proper examination, I readily perceived an hard body confined in the interior membrane of the intestine. With the help of a pair of forceps, I
extracted

extracted two portions of a brittle black substance; which, on careful inspection, appeared to be bread toasted nearly to a cinder: the two pieces, which were whole before the extraction was attempted, might be together about an inch in length, half an inch in width, and one third of an inch in diameter.

The patient remembered to have swallowed something with considerable difficulty two days before, while partaking of some soup; which was probably the same morsel of bread that distressed him upon this occasion.

Does it not appear from this case, that bread when toasted is less fit for digestion than some persons would have us believe; and that it affords but little nourishment compared with that which is moderately baked?

However trifling the circumstances of the above case may be regarded in its earliest stage; there can be no doubt entertained of the probability of its terminating very seriously, if the patient had not applied for speedy relief: inflammation, abscess, and all their consequences, might have ensued, if the efforts of nature, or the power of aperient and antiphlogistic remedies had alone been trusted to.

In similar instances, without losing time by
endeavours

endeavours to relieve the patient's sufferings by medicine, it will be immediately proper to subject him to a careful examination. If the simple introduction of a finger be insufficient to disengage the extraneous body, and it can be felt adhering to the *rugæ*, or piercing the coats of the rectum, a pair of blunt-pointed scissars, or forceps, (as the case may indicate) should be gently conducted upon the finger, in order to divide, break in pieces, or loosen the foreign substance: if a pointed bone, or other hard and sharp body, should be confined across the gut, endangering the neighbouring parts, it will be prudent to empty the urinary bladder, previous to any attempt to remove it by mechanical means: and, should the pain, and other ill effects become urgent, it might be necessary, after milder methods had proved ineffectual, to make a judicious incision either into the rectum, or circumjacent integuments, as the peculiarities of the case should require to facilitate the extraction. To obviate the inflammation, and its concomitant symptoms, leeches, anodyne and laxative clysters, with the usual antiphlogistic remedies, ought to be diligently employed.

Instances of the kind above related, with
 VOL. VI. I suitable

fuitable remarks, are recorded by several practical authors ; but the reader may spare himself the trouble of perusing some of them, by consulting the *Memoires de l'Académie Royale de Chirurgie*, Tom. 1. p. 540, et seq. 4to Edit.

Newman Street, Oct. 6, 1794.

VII. *A Case of Aneurism of the Crural Artery ; communicated in a Letter to Dr. Simmons, by Mr. Thompson Forster, Surgeon on the Staff of the Army, and Surgeon to Guy's Hospital.*

TO DR. SIMMONS.

Dear Sir,

TO the two cases of Aneurism which you have done me the honour to insert in the fifth volume of *Medical Facts and Observations*, I am desirous of adding the following, as I flatter myself it will tend still further to elucidate the peculiar utility and advantages of the operation in question.

Believe me, Dear Sir,

Your's, &c.

Nov. 3, 1794.

THOMPSON FORSTER.

CASE.

C A S E.

Lawrence M'Carthy, a labouring man, aged thirty-seven years, was admitted, as my patient, into Guy's Hospital, on the 30th of July 1794, for the cure of an aneurism of the crural artery.

About nine months before his admission, he had perceived a small tumor on his right thigh; near that part where the crural artery dips under the triceps muscle; as it occasioned no inconvenience, nor prevented his working, he took but little notice of it; it came spontaneously, without any external violence, and remained stationary for near six months before it became painful: when the tumor had acquired the size of an egg, a pulsation was perceptible in it, but not before.

At this period of the disease he was advised to foment the part, and to make use of liniments: this he continued to do for some time; but finding no relief from these remedies, he applied to a surgeon, who recommended the use of a bandage, which he made use of for near three months, but without any abatement of the pain; and the tumor in the mean time had increased to a very considerable

size, and the limb in general had acquired something more than its natural bulk.

The patient, naturally hypochondriacal, became anxious, irritable, and dejected; complaining of great pain in the limb, and particularly in the tumor, which was in some measure eased by pressure. In this state he came into the hospital; and his general habit having been lowered by bleeding, purgatives, and a suitable regimen previously to the operation, I performed it on Monday, the 11th of August, by making an incision in the course of the lower edge of the sartorius muscle, and about an inch below where the profunda is usually given off. Having laid bare the artery, * I passed a ligature under it with an eyed probe, and applying the stick, surrounded by adhesive plaster, &c. as described in the former cases †, the ar-

* With a view of conveying to the reader a more precise idea of the operation, I have made a sketch of the parts concerned in it, from a subject dissected for the purpose. See the annexed engraving (plate 1, fig. 2.) in which *a* refers to Poupart's ligament; *b* to the crural artery, with a ligature passed under it at the part where it was tied; *c* to the profunda; and *d* to the sartorius muscle. It seems hardly necessary to remind the reader that the object of this sketch being merely to point out the seat of the operation, the parts are delineated in their natural state.

† Vide Vol. V. p. 6.

tery

tery was thus surrounded, and by these means equally compressed; the pulsation below of course ceased: but, for fear of a sudden hæmorrhage, I passed a second ligature about half an inch above the former, laying it loose, that an assistant might instantly tie it in case of such an accident.

August 21st. The first ligature, with the stick, came away with ease.

August 22d. The second ligature came away with equal ease.

An account of the state of the pulse at the wrist, and of the temperature of both limbs, at the ham, and at the foot, was taken every day with great accuracy by Mr. G. Babington, according to the annexed Table *, until August the 27th, when the temperature of each was found to be equal.

The size of the tumor gradually decreased, and the patient, having the perfect use of his limb, was dismissed, cured, October 10, 1794.

The preceding case differs materially from the two former, not only in the circumstance of the tumor in this having been situated in the

* See page 119.

upper part of the thigh, so that the artery could not be secured lower than about an inch below where the profunda is usually given off, but likewise in the very great pain the patient endured both night and day for three weeks before the operation. The tumor was as considerable, but the enlargement of the limb below it was much less than in the former cases. After the operation, the symptoms were much slighter than in the other cases, probably owing to the low state I thought it proper to reduce the patient to for the purpose; and the ligature came away on the tenth day after the operation without the least trouble. But the circumstance in which it differed the most essentially from the other two, was, that the tumor was completely absorbed in seven weeks, and the patient had then acquired a perfect use of the limb, while, in the former cases, the patients did indeed acquire the use of their limbs, but the tumors, though lessened and free from pulsation, still remained.

TABLE.

TABLE.

Day of the Month.	pulse at wrist	tem. of arm.	tem. right ham.	right foot.	left ham.	left foot.	Time of day when the obs. were made.
August 11		68 $\frac{1}{2}$ °	98 $\frac{1}{2}$	94 °	97 °	96 °	10 $\frac{1}{2}$ P. M.
12		68 $\frac{1}{2}$	97	91	91	89	8 $\frac{1}{2}$ A. M.
	128	70	99	91	94	93	10 $\frac{1}{2}$ P. M.
13	109	68	98	92	90	88	8 $\frac{1}{2}$ A. M.
	112	71	100	95	98	95	10 $\frac{1}{2}$ P. M.
14	104	68	98	91	91	91 $\frac{1}{2}$	8 $\frac{1}{2}$ A. M.
	116	72	99	96	96	96	8 $\frac{1}{2}$ P. M.
15	96	69	97	91	94	88	8 $\frac{1}{2}$ A. M.
	112	72 $\frac{1}{2}$	97	93	94	95	8 P. M.
16	97	72	98	93 $\frac{1}{2}$	94	90	8 A. M.
	112	73	98	95	94	94	8 P. M.
17	96	71	98	92	95	89	8 A. M.
	112	74	97	94	94	94	8 P. M.
18	92	70	93	91	92 $\frac{1}{2}$	89	8 A. M.
	110	72	97	92	94	93	9 P. M.
19	100	68	94	90	92	91	8 A. M.
	124	71 $\frac{1}{2}$	101	96 $\frac{1}{2}$	97	97	8 $\frac{1}{2}$ P. M.
20	114	67	100	93	96	94	8 $\frac{1}{2}$ A. M.
	116	70	99	95	95	94	8 $\frac{1}{2}$ P. M.
21 First ligature and stick came away with ease, there being a perfect solution of continuity.							
	100	66 °	97 °	8 °	93 °	86 °	8 A. M.
	100	69	98	92	95	94	8 P. M.
22 Second ligature was removed.							
	100	69 °	96 °	86 °	93 °	84 °	8 $\frac{1}{2}$ A. M.
	108	69	98	94	97	95	9 P. M.
23	100	67 $\frac{1}{2}$	96	91	93	90	9 A. M.
	104	70 $\frac{1}{2}$	98	94	95	93	8 $\frac{1}{2}$ P. M.
24	96	69	97	89	95	87	10 A. M.
	104	69	99	95	98	95	8 P. M.
25	104	66 $\frac{1}{2}$	98	93	96	92	8 $\frac{1}{2}$ A. M.
	106	69 $\frac{1}{2}$	95	92	93	91	8 P. M.
26	100	64	96	90	91	86	8 A. M.
	106	66	98	93	94	90	8 $\frac{1}{2}$ P. M.
27	100	64	96	91	92	91	8 $\frac{1}{2}$ A. M.
	95	63 $\frac{1}{2}$	96	90	96	90	8 P. M.

VIII. *An Account of a Key Instrument of a new Construction ; with Observations on the Principles on which it acts, in the Extraction of Teeth, and on the Mode of applying it. By Mr. Robert Clarke, Surgeon at Sunderland, in the County of Durham. Communicated in a Letter to Mr. Anthony Carlisle, Surgeon of the Westminster Hospital, and Reader of Anatomy in London ; and by him to Dr. Simmons.*

To Mr. CARLISLE.

SIR,

WITH this I send you a Key Instrument, for the Extraction of Teeth, of a construction different from any in common use, and which in practice fully answers to the expectations I had formed, *a priori*, from a careful examination of the principles of its action.

I cannot, perhaps, give you a clearer idea of its advantages, than that which you will obtain by pursuing the same train of investigation which I followed myself. I shall therefore proceed to lay it before you, that I may more thoroughly convince you of the propriety of the alteration I have made, or be corrected by your pointing out any error I may have fallen into.

IN

Fig. VII.




Fig. VIII.




Fig. IX.





Fig. X.



Key Instr. of

Fig.

Key Instr. of

Fig.

Key Instr. of the Comm. inside of the Jaw

Fig.

Key Instr. of the old the inside of the Jaw

struction

n fixed on the

n fixed on

In the first place then, it appeared to me that as the fulcrum, or point, upon which the tooth is carried round as on a center, is that part of the bolster which rests upon the gums, the axis of motion of the instrument would always be found by drawing a line through that point and the middle of the handle; and consequently that the old construction of the Key Instrument was free from an inconvenience which attends the more modern one: I mean the axis of the bolster and axis of the shank making an angle with each other; on which account it is disposed to shift its point of action on the gums, and to raise the tooth in a plane inclined to the throat, instead of a vertical one, as may be clearly seen by inspecting Figures I. II. (Plate II.*) where *a, b*, represent the axis of motion; *c, d*, the direction in which each instrument raises the tooth; and *e. f.* (Fig. II.) the axis of the bolster.

Now as the line of direction in Fig. I. is perpendicular to the jaw, it is needless to say that it is highly preferable to Fig. II. where the line of direction is inclined backward, making the

*. It seems right to observe here, that all the figures of this plate are on a reduced scale of two thirds of their proper size.

extraction of the tooth more difficult, and exposing that which is situated behind it to be driven from its socket, or even to be caught in the arch of the claw. Besides this, the bolster rests only upon the corner *d*, adding greatly to the injury of the gums.

The construction then of the Key-instrument delineated in Fig. I. would seem perfect, were it not that in drawing teeth inwards, with respect to the jaw, the fore teeth prevent its due application, confining it to the direction represented in Figure V.

To remedy this imperfection I have made the instrument with a bend in its shank, to clear the fore teeth, and to allow its proper application, as in Figure III. where the same observations and references apply as in Figure I. and therefore it is unnecessary to repeat them. But in order that the comparative merits of the three instruments may be seen at a glance, I have added Figures IV. and V. wherein the axis of motion, and the direction of the rising tooth, are shown by dotted lines.

Having fully considered what relates to the direction of the tooth, I shall next examine the mechanism which takes hold of it. For this purpose recourse must be had to the engraving.

Let

Let *a, b, c*, Figure VI. represent an end view of a Key instrument, fixed upon a piece of hard, smooth wood. Then it is obvious, that if it be turned from left to right, by means of its handle, it will break the wood in the direction *d, c*, and cause the upper fragment to revolve on the point *c*, as a center. It is equally obvious, that if a line be drawn from the point *a*, crossing the opposite surface of the solid *e, f*, at right angles, the counterpoise of the claw will fall into that line before it can take hold; for then the point *b*, is at the greatest possible distance from the surface *e, f*; consequently if the instrument be placed as in Figure VII. the point *c* will descend; or, if as in Figure VIII. it will ascend until it coincides with the line *a, b*.

I shall now endeavour to apply this to practice. Let 1, 2, 3, in Figure IX. represent a tooth with its roots fixed in a section of the jaw, and its corona engaged in a Key-instrument; then it will readily appear that upon the action of the instrument, the tooth will be drawn from its socket, and carried round the point *b*, as a center, rather than the joint substance of the tooth and jaw be broken in the line *a, b*, as happens in Figure VI. This however happens

only under particular circumstances : For if the bolster be placed too high, the tooth will be broken ; and if too low, the alveolar process will always be torn away with it. It is therefore a matter of importance to determine the best point of contact for the bolster, and this I have uniformly found to be at two-thirds the depth of the tooth, the claw being fixed at one third, as represented in Figure IX.

It will always be easy to ascertain this point, by attending to the size of the corona, and the part of the jaw where the tooth is situated ; and equally so to make the instrument act upon it, by using a larger or smaller claw as the case may require. For illustration, however, I shall refer to Figure X. which represents a piece of wood grasped by the tooth instrument in the same manner as in Figure VI. Now if a larger claw, shewn by the dotted line, be used, the bolster will fix higher upon the wood than before. For as the center pin of the claw will always rest in the line *a, b*, the bolster must rise higher before it can come into contact. But notwithstanding the use of a larger or smaller claw, in proportion to the size of the tooth, enables us to fix it at a proper height, the use of a very disproportionate one is always inconvenient,

inconvenient, by depriving us of the use of the crank, in drawing teeth inwards, and by encroaching upon the cheeks in drawing them outwards. I have therefore in the construction of this instrument, taken care to make the bolster of such a depth, as to be free from either inconvenience.

The form of the bolster is by no means a matter of indifference; for if it be too small, it presents so small a surface to the gums, that the pressure made upon them, by the extraction of a tooth moderately firm, cuts them through, and even penetrates the bone, especially if the bolster be of the usual form. I have therefore been careful to make it of a proper size, and to give it a prolate spheroidal figure, as being the least disposed to injure the gums, and applicable with exactness and ease to all parts of the mouth; and in order still further to guard against this bruising of the gums, I wrap the bolster to the thickness of a line, with tow, wound on as tight as I can, before I slide forward the bolt and put in the claw.

I have also been attentive to the form of the claws, that they may touch the tooth only with their points. And the instrument is so contrived, that they can be quickly changed or
turned

turned to an opposite direction as the case may require : this is done by means of a sliding bolt, instead of a screw, which passes through the claws.

I have always found that when the tooth is to be turned from right to left in drawing it, that the handle answers best placed perpendicularly ; and when from left to right, horizontally. The reason of this will be obvious, if we consider that in the first case, the pronator muscles of the operator's arm, which are those exerting the force, act with most advantage when the hand is vertical ; and in the second case, that the supinators act most advantageously with the hand prone. I have therefore contrived the handle so that it may be easily turned, as often as there is occasion to turn the claw.

I am, Sir, &c.

Sunderland,

ROBERT CLARKE.

Aug. 18, 1794.

IX. *An Account of a new Species of Swietenia (Mahogany); and of Experiments and Observations on its Bark, made with a View to ascertain its Powers, and to compare them with those of Peruvian Bark, for which it is proposed as a Substitute: Being an Abstract of a Paper on this Subject, addressed to the Honourable Court of Directors of the United East-India Company. By William Roxburgh, M.D.*

THE species of *Swietenia* described in this paper, and which Dr. Roxburgh names *Swietenia Febrifuga* *, is a native of the mountainous part of the Rajamundry Circar, North of Samulcotah and Peddapore. It is a very

* Dr. Andrew Duncan, junior, who has made this new species of *Swietenia* the subject of a very ingenious inaugural Dissertation, gives a good reason for preferring, as a trivial name, the Hindoo appellation, *Soymida*, to one founded on its medicinal properties; similar properties, he observes, being ascribed by Dr. Wright (London Medical Journal, Vol. VIII. p. 286) to the mahogany tree of Jamaica (*Swietenia Mahagoni*), another species of the same genus.—Vide *Tentamen inaugurale de Swietenia Soymida*; Auctore Andrea Duncan. 8vo. Edinburgi, 1794. EDITOR.

large

large tree, known among the Hindoos by the name of *Soymda*, and flowers about the end of the cold or beginning of the hot season. Its seeds ripen in three or four months after.

Of this tree Dr. Roxburgh gives the following botanic description :

“ TRUNK. Very straight, rising to a great height, of a great thickness, and covered with a grey, scabrous, cracked bark.

“ BRANCHES. Numerous, the lower spreading, the higher ascending, forming a very large shady head.

“ LEAVES. Alternate, about the extremities of the brachlets (*turiones*) abruptly feathered, about twelve inches long.

“ LEAFLETS. Opposite, very short, petiolated, three or four pair, oval, obtuse, or end-nicked, the lower side generally extending a little further down on the petiolet than the upper; smooth, shining; from three to five inches long, and from two to three broad, the inferior smallest.

“ PETIOLE. Round, smooth, about nine to ten inches long.

“ STIPULES none.

“ PANICLE. Very large, terminal, diffuse,

“ fuse, bearing great numbers of middle-sized,
 “ white, inodorous flowers.

“ PEDUNCLE and PEDICLES. Round
 “ and smooth.

“ BRACTS. Very minute.

“ CALYX. Below, five-leaved; LEAF.

“ LETS. Oval, deciduous.

“ COROL. Petals five, inverse, egged,
 “ obtuse, concave, expanding. NECTARY.
 “ Not quite half the length of the petals, a
 “ little bellied; mouth ten-toothed, teeth bi-
 “ fid (two-cleft).

“ STAMEN. Filaments ten, very short,
 “ inserted just within the mouth of the nectary.

“ ANTHERS. Oval.

“ PISTIL. Germ conical. STYLE. Thick,
 “ tapering. STIGMA. Large, targetted, shut-
 “ ting up the mouth of the nectary.

“ PERICARP. Capsule egged, large, five-
 “ celled, five-valved, valvelets gaping from
 “ the top.

“ RECEPTACLE. In the centre, large,
 “ spongy, five-angled; angles sharp and con-
 “ nected, with the sutures of the capsule, be-
 “ tween them, deeply fulcated.

“ SEEDS. Many in each cell, imbricated,
 “ obliquely wedge-shaped, enlarged by a long

“ membranaceous wing, inserted, at the upper
 “ point of the wing, into a long brown speck
 “ on the upper part of the excavations of the
 “ receptacle : all the rest of the receptacle is
 “ white.”

The wood of this tree, we are told, is of a dull red colour, remarkably hard and heavy ; and is reckoned, by the natives, by far the most durable timber they know ; on which account it is used for all the wood work in their temples.

The bark of the trunk and large branches, of large and middle-sized trees, is covered with a dark rusty-coloured coat, of about an eighth of an inch in thickness, which cracks in various directions, and sometimes peels off in irregular pieces, according to the directions of the cracks. Immediately under this is a very firm, but brittle coat, of about three-eighths of an inch in thickness. When first cut, it is light-coloured ; but on drying, or even exposure to the air for a few minutes, it turns to a reddish brown. The inner lamina are thin, consisting of tough, lighter-coloured layers.

The bark of the younger branches is not cracked, is pretty smooth, of a much lighter colour,

colour, and has not the rusty coat above described, but has often many blotches of various coloured lichen over it: it consists wholly of the brown, solid, and inner layers.

The outer rust-coloured layer of the trunk has but little taste; the other two possess a little aromatic smell, and their taste is very bitter and astringent, accompanied with something aromatic, but in a trifling degree. There is nothing disagreeable in the taste, more than may be expected from a pure, simple, strong bitter and astringent united. The middle lamina are easily reduced to a very fine rose or light brown-coloured powder.

Cold water, in the course of an hour, our author observes, acquired from this bark a deep but clear reddish colour. The most minute portion of a chalybeate (one drop of a solution of twenty grains of sal martis in an ounce of water) instantly changed a much-diluted cold infusion to a deep purple, which, on standing, became darker and darker, with a reddish tinge; and no decomposition took place till about the tenth day; the colouring matter then began to separate, and fall to the bottom in black flakes, leaving the liquor almost colourless. If the infusion was some days (from four to thirty)

old, the colour produced by the martial solution was as instantaneous as when fresh, and deeper; and at no period, up to thirty days, did it show the least tinge of green. Ten times the same quantity of the same martial solution, it seems, did not produce so great a change upon a similar infusion of the common pale Peruvian bark; and its effect on the latter was much slower. Its bitter qualities are also described as much more intense than those of the common sort of Peruvian bark.

The infusion, we are told, bears to be mixed in any proportion with spirits, without becoming turbid, or producing any kind of decomposition. The first decoction is considerably deeper-coloured than the infusion (which colour it retains in passing the filter), and possesses the same powers in a higher degree, but does not retain them so long, nor is it so pleasant to the taste. On standing any length of time with the chalybeate, the colour becomes pale, and is sooner decomposed than the cold infusion: on standing some days it lets fall a small quantity of a reddish, earthy fecula, which is intensely bitter and astringent; the superincumbent liquor becoming gradually clearer and clearer, and at the same time of a deeper red, much resembling the tincture. Lime-water added to the decoction,
infusion,

infusion, or diluted tincture, darkened them considerably, and caused in all a copious deposition of reddish brown fecula, which became purple coloured in twenty four hours. The decoction, it is observed, gave the greatest quantity of fecula. An infusion of pale Peruvian bark, prepared in every respect the same as the infusion of Swietenia bark, was treated with lime-water in the same manner, and formed a separation, but in a much less degree.

Bark of *Melia Azadirachta* (Margosa tree) treated exactly in the same manner, formed a separation of a lighter-coloured fecula, in a much greater quantity than the common Peruvian bark, but much less than the Swietenia bark.

The clear reddish-coloured liquor, we are told, that floats over the precipitate caused by the addition of lime water, is void of astringency to the taste, or has it only in a trifling degree; but for a farther proof, it seems, a chalybeate was employed, which did not in the least darken this liquor; but a greenish tinge was produced, together with a further decomposition and precipitation of a reddish fecula. This experiment, our author thinks, serves to show that at least the astringent part of the bark is entirely thrown down by lime-water; and he considered this as so interest-

ing a point, that he repeated the same experiment with this, as well as with other astringent barks, and found the result exactly the same.

The same chalybeate added to lime-water of the same strength as that employed in the above-mentioned experiments, produced a small, green cloud; the Swietenia bark infusion thrown into this produced a muddiness, and soon after, a copious precipitation of dirty-coloured fecula.

An infusion of this bark in lime-water is deeper coloured than the plain infusion, but possesses very little bitterness, and still less astringency. A chalybeate added to this infusion rendered its red colour a little deeper only, and no decomposition took place: after standing some time, the infusion had no taste of the lime-water.

From these experiments, Dr. Roxburgh considers lime-water as a very improper addition; and observes that, in this respect, they agree with those made by Dr. Irving on the red and quilled Peruvian barks.

Vitriolic acid rendered the first decoction, or watery infusion, paler; and, upon standing, it became a little turbid, and let fall a small quantity of a light-brown sediment.

Vinegar had the same effect,

Mild, or caustic vegetable alkali, or mild
fossil

fossil alkali, soon deepened and rendered brighter the cold watery infusion or decoction, nor did any decomposition take place in forty-eight hours.

Mild magnesia, simply added, rendered the colour of the infusion paler, without sensibly altering the taste.

Alum has been at times successfully employed for the cure of intermitting fevers, and the analogy it bears to other tonics renders it a likely remedy. Our author was therefore desirous to try what would take place on adding it in a small quantity to infusions and decoctions of this bark. The addition, it seems, rendered their colour paler, and a little decomposition took place, with a precipitation of a small quantity of a light-brown fecula: to the taste it increased the astringency without sensibly diminishing the bitter; but with alum they did not change their colour when a solution of green vitriol was added.

Eight ounces of the coarse powder were boiled in ten pints of soft well-water to four pints; the residuum was repeatedly boiled in fresh parcels of water, exactly in the same manner for eleven times, when the liquor

came off still much coloured, but tasteless, and showed no signs of astringency with the chalybeate; the tenth decoction excepted, which did show signs of astringency, as it was darkened a little by it:

The fresh decoction of common Peruvian bark, made similarly, but in a smaller quantity, struck slowly about as deep a colour with the same chalybeate, as the fourth or fifth decoction of Swietenia bark did quickly.

As the eleventh decoction was tasteless, although coloured, it was thrown away; the other ten had been regularly strained, while hot, and suffered to stand till perfectly cold, then poured off, clear from sediment; they were mixed, and evaporated to a hard extract, which weighed two ounces and three-quarters. The extract, when soft, was of a dark red colour, flavourless, smooth, homogeneous, and unctuous when rubbed between the fingers and thumb. The taste of the decoction was well preserved in this extract; the most minute part of it, dissolved in water, struck a black colour with martial solution as quickly and as deep as the decoction itself, but the taste was not so strong as might be expected from that of the bark. This, our author thinks,
might

might perhaps be owing to the more fixed, inert parts, extracted by the long and repeated boilings (which lasted two days) being mixed in the mass of extract. But this, he observes, would not be the case, or but in a small degree, with one prepared from only one or two boilings. To determine this point, he boiled one ounce of the powdered bark in two pints of water, pretty briskly, down to one pint; after the liquor was poured off, to the residuum were added two other pints of water, and boiled in the same manner. The decoctions were mixed, and evaporated to a dry extract, which weighed two drachms and a half, and was in taste, &c. much as the former from ten coctions; the proportion of extract from two boilings is therefore, he observes, nearly equal to that of ten: so that, although the decoctions were highly coloured, and considerably bitter and astringent, even to the tenth, yet they could have contained but a small portion of the powerful qualities of the bark.

The residuum, when perfectly dry, weighed four drachms and a half; and spirit of wine being poured on it, though assisted at times with the heat of the sun for many days, extracted
neither

neither colour nor taste, so completely had the virtues of the bark been extracted by the water.

Dr. Roxburgh observes that the dry extract imbibes much moisture when the weather is damp; so much as to make it stain the fingers, or any thing that touches it: that it melts readily in the mouth; is easily soluble in water and in spirits; and, like the decoction and tincture, bears to be mixed without decomposition. These solutions and mixtures, we are told, resemble much the original decoction and tincture, and their mixtures, both in taste and colour.

Should this ever become the valuable drug it promises, it would be advisable, our author thinks, to have the extract prepared on or near the spot where the trees grow. If this is done during the hot season, the evaporation, he observes, might be effected by the heat of the sun and hot winds, which would certainly produce a much more elegant, efficacious extract than could possibly be prepared in any other way or place, and would also preclude every idea or chance of its being sophisticated.

This bark, he finds, contains much mucilaginous matter, the cloth that the decoctions
were

were strained through, having become, when dry, stiff as if starched. This, he thinks, may account for the decoctions remaining so many days turbid, which is, no doubt, he adds, favourable for the action of the stomach upon the bark. The late Dr. Fothergill, he observes, recommended an addition of some mucilage to decoctions of common bark, in order to keep them turbid, that the active parts might be kept more completely suspended in the liquor *.

In the way of distillation, this bark, it seems, yields nothing, not the smallest apparent quality, either with water or spirits. In this respect, Dr. Roxburgh thinks, it resembles exactly both the pale and red Peruvian barks, viz. in having its powers or virtues of a very fixed nature.

Rectified spirit of wine extracts from the bark a clear, deep red tincture, possessing the astringency of the watery infusion or decoction, and more of the bitter. If not too strong, it makes, we are told, one of the most pleasant bitters we are in possession of; and it bears to be diluted with water in any proportion, without decom-

* Med. Obs. and Inq. Vol. I. p. 321. 2d Edit. 8vo. London, 1758.

position,

position, which renders it in many cases the more desirable.

Four ounces of powdered bark were infused, by our author, for eight days, in three pints of French brandy; these were poured off, and four pints more of the same brandy added, which, after standing four days, were also poured off: both these infusions were mixed, and he drew off, by distillation, a quantity of the spirit, which (as before observed) did not in the least partake of any of the qualities of the bark: the rest was gently evaporated to a dry extract, which weighed nine drachms. The extract itself was of a much darker colour than that procured by water, and was dried with more difficulty; but the taste of the two extracts was much the same. The residuum was boiled in six pints of water to two, and the decoction was found to be still pretty strong to the taste, both in bitterness and astringency. This induced him to repeat the boiling, twice more, with fresh parcels of water; and the last decoction, though weak, was still bitter, and showed signs of astringency, with a martial solution. These four decoctions were mixed and evaporated to a dry extract, weighing three drachms, which added to the spirituous extract,

tract, made in all twelve drachms, from four ounces of powdered bark, and agreed nearly with the quantity procured by water alone.

The antiseptic powers of this bark, according to our author's experiments, are not inferior to its bitter and astringent qualities; for watery infusions in open phials kept perfectly good for sixty days, without any tendency to fermentation, except a few air bubbles, which they discharged about the second day; indeed they acquired strength, we are told, as the colour produced at the end of that time (sixty days), by the addition of a chalybeate, was darker, and as instantaneous as at any prior period.

Sixty grains of the lean of raw mutton were preserved sweeter and firmer in an infusion of ten grains of this bark in four ounces of water, than an equal quantity of the same mutton in a similar infusion of pale quilled Peruvian bark. The flesh was tinged red by the infusion of Swietenia bark, and its fibres were firm and distinct at the end of twelve days; while that preserved in the Peruvian infusion was white, and its fibres softer, and infinitely more fetid.

Almost all the foregoing experiments, it is
observed,

observed, were made first with bark of the smaller branches, and again with bark of the trunk of a large tree; the latter was evidently strongest.

The seeds of this tree are described as a strong, simple, pleasant bitter, without any of the astringent power. The leaves possess nearly if not all the astringency of the bark, and a very large proportion of its bitter; but their taste is said to be not so agreeable either in substance or in infusion.

From the foregoing analysis, Dr. Roxburgh ventures to draw the following conclusions:

First. That the active parts of the bark of this species of *Swietenia* are much more soluble than those of Peruvian bark, particularly in watery menstrua.

Secondly. That it contains a much larger proportion of active (bitter and astringent) powers, than Peruvian bark.

Thirdly. That the watery preparations of this bark remain good much longer than similar preparations of Peruvian bark.

Fourthly. That the spirituous and watery preparations bear being mixed in any proportion, without decomposition.

Fifthly. That the bark in powder, and its preparations,

preparations, are much more antiseptic than Peruvian bark, or similar preparations of it.

Now, since this bark yields so much of its virtues to cold water, as to preserve flesh from corruption, in a hot climate, with the thermometer from 87° to 102° , it is reasonable, he contends, to suppose it will yield still more of its tonic and antiseptic virtues in the stomach, where it meets with the most powerful solvents : we have therefore, he thinks, much to expect from it in the cure of gangrene and other putrid diseases.

Bitters and astringents, in a separate state, our author observes, are considered as tonic remedies ; but when found combined in the same substance, they become still more powerful : it is from these qualities, he contends, that the best judges allow the Peruvian bark to derive its virtues. On this point he quotes the authority of Dr. Cullen, who has remarked, “ that the
 “ recurrence of the paroxysms, in intermitting
 “ and remitting fevers, depends on the recurrence of atony in the extremities of the arterial
 “ system; hence they are prevented by such
 “ tonic medicines as obviate this atony : a
 “ great variety of astringents and simple bitters
 “ have been found to answer that end, but
 “ none, hitherto discovered, so effectually as the
 “ Peruvian

“ Peruvian bark, on account, it is thought, of
 “ its possessing those powers conjoined *.”

The antiseptic qualities of Peruvian bark, our author observes, are also great; hence its use in the cure of all febrile putrescent disorders, attended with debility, putrid ulcers, &c.

From the evident qualities of this new bark, and from the successful experience he has had with it, in intermittent fevers†, &c. Dr. Roxburgh

* Treatise on the Materia Medica.

* Histories of several of these cases have been communicated by Dr. Roxburgh to the College of Physicians at Edinburgh, and an account of them is given by Dr. Duncan in the dissertation referred to in a former note, together with the results of several trials made with this bark, by his father, in the Clinical Ward of the Royal Infirmary at Edinburgh. We shall take the liberty of transcribing this part of his work :

“ Morbus, quo Roxburgius hunc corticem sæpissimè adhi-
 “ bendum curavit, febris quotidiana apud Cullenum nun-
 “ cupatur. Rariùs ex toto, sed ex partè, et ad breve tantum-
 “ modo tempus, remittens, periculosissimus erat. Ægroti ferè
 “ omnes hoc morbo correpti fuerant, dum incolabant istos
 “ montes ingentes, qui Indiæ peninsulam transcurreunt. Inter
 “ hos montes sylvæ opacæ, densa ferarum tectæ, convalles pa-
 “ ludosæ, hominum generi pestíferas, ubique obumbrant. Se-
 “ des est indigenis etiam, consuetudine licet obfirmatis, insa-
 “ lubris, advenis autem adeo perniciofa ut pauci, perpauci
 “ quidem, quos dira necessitas inter hos montes hiemare coege-
 “ rit, morbo hoc atrocissimo immunes sint. Tali febre, tali
 “ tempestate

burgh has every reason to imagine it will prove equal, if not superior, to the Peruvian bark, for every purpose for which that medicine is used.

Our

“ tempestate laborantium ne dimidiam quidem partem con-
“ valescere Roxburgius affirmat.

“ Cal. Junii, A. D. 1791. Indus annos natus viginti,
“ habitus tenuis, nonnullis antè mensibus, dum prope montes
“ occupabatur, febre quotidianâ affectus erat. Corticem
“ Cinchonæ officinalis aliquantisper sine fructu assumpserat;
“ idcirco Roxburgius, et quia ipse parvas corticis Soymidæ
“ quantitates impunè adhibuerat, ægro nihil à periculo ab-
“ horrenti grana viginti pulveris ex aquæ cyatho sumenda
“ præscripsit. Duabus exinde hōris, scrupuli duo adhibiti
“ sunt; et, post simile temporis intervallum, drachma. Cor-
“ tex ægro nequaquam ingratus erat, alvumque solvit. Æger,
“ cortice postea ad drachmam, unaquâque intermissione, as-
“ sumpto, triduo febre immunis erat.

“ Pridie Iduūm Augusti, A. D. 1791, J— V— Lusit-
“ tanus*, annum agens quadragesimum quintum, ejusque duæ
“ filiæ, altera sex, altera tres annos nata, manserant aliquan-
“ diu, inter mensem proximè præteritum, intra montium ter-
“ minos; initioque mensis labentis, febre quotidianâ, quæ
“ nihil fermè quicquam remisit, affecti sunt. Febre remit-
“ tente, semper alterâ quaque horâ fumebant aquæ ex cortice
“ Soymidæ* pater sescunciam, filia major natu unciam, et
“ minor semunciam. Duos post dies, à morbo valebant.

* “ R. pulv. cort. Swiet. Soymidæ unciam unam,

“ aquæ fontanæ libras duas,

“ Misceantur, et phialâ prius agitâtâ, modo præscripto sumantur.”

* Vide p. 148.

VOL. VI,

L

“ Morbus,

Our author next enumerates different species of Cinchona, viz.

First.

“ Morbus, quo hi quatuor ægroti laborabant, partim ob
 “ anni tempus, quo febre correpti sunt, atque partim ob
 “ tempestatis ficcitatē, solito levior erat; atque Roxbur-
 “ gius, propter ægrorum debilitatē, neque evacuantia ad-
 “ hibebat, nec intermissiones expectabat.

“ xv. Cal. Sept. A. D. 1791, B— Lusitana, habitûs
 “ infirmi, nonnullos dies, febre gravi, nunquam ex toto re-
 “ mittente, laboraverat. Antimonium tartarifatum ex multâ
 “ aquâ, partitis vicibus, usque ad vomitionem, adhibuit.
 “ Postero die drachma corticis Soymidæ, in remissione mi-
 “ nimè adhuc notabili, ter assumpta est. Intermissio proxima
 “ plenior evasit, atque, ex corticis usu, biduo postea morbus
 “ ipse, simulque diarrhœa quâ laboraverat ægra, cessârunt.

“ Menfe Septembris, A. D. 1791, J. E— decurio Euro-
 “ pæus, annos natus quadraginta, febre remittente graviter
 “ affectus est. Recessus principio ferè nulli, ex usu præpara-
 “ torum ex antimonio notabiliores evaserunt; et æger, quan-
 “ quam omni generi intemperantiæ deditus, cortice ter sin-
 “ gulis intermissionibus adhibito, paucis diebus convaluit.

“ vii. Cal. Sept. A. D. 1791, T. L— annos natus
 “ octodecim, quosdam dies febre biliosâ laboraverat; cujus
 “ recessus, etiam post antimonii tartarifati usum, parum
 “ notabiles erant. Debilitate autem urgente, scrupuli duo
 “ corticis Soymidæ, omni recessu, ter adhibebantur, et, ad
 “ alvum solvendam, lixiva tartarifata.

“ A cortice autem nihil proficiente, iiii. Cal. decessum
 “ est; atque medicamentis idoneis assumptis, febris prorsus
 “ fere,

First. *Cinchona Officinalis panicula brachiata*;
to this species, he observes, belong the pale,
quilled,

“ fere, statis temporibus, intermisit. Soymida nunc iterum
“ adhibita, quatuordecim diebus, morbum penitus fugavit.

“ Menſe Septembris, A. D. 1791. S— nutrix lactans,
“ annos nata triginta quinque, febre quotidianâ correpta
“ eſt. Alvo, inter primam intermiſſionem, ſodâ vitriolatâ
“ ſolutâ, morbus triduo cortice Soymidæ depulſus eſt; ſed
“ lac interim fluere ceſſaverat.

“ Menſe Septembris, A. D. 1791. Indus, ſervus domeſ-
“ ticus†, febre ſingulariter intermittente ægrotavit. Sub
“ occaſum ſolis, acceſſit febris gravis, quæ horâ nonâ veſ-
“ pertinâ intermiſit. Oriente autem ſole, iterum acceſſit,
“ atque, horam circiter nonam ante meridiem, denuo inter-
“ mittens, ægrum viribus integrum reliquit. Exinde cor-
“ tice Soymidæ ter, ſingulis intermiſſionibus matutinis, ad
“ ſcrupulos duos adhibito, triduo morbus omnino evanuit.

“ J— R— Europæus annum agens trigeſimum, vitio pul-
“ monis multum debilitatus, incunte Octobri febre quotidi-
“ anâ, cui erant acceſſiones veſpertinae, affectus eſt. Tertiâ
“ intermiſſione, duo corticis ſcrupuli bis adhibiti alvum
“ magnopere ſolverunt. Soymidâ nihilominus continuatâ,
“ æger quatuor diebus à febre valebat.

“ Pridie Iduûm Decemb. R— miles Indicus, annos natus
“ triginta, febre quotidianâ tredecim dies, medicamentis
“ vernaculis nihil proficientibus, laboraverat. Intermiſſione
“ proximâ duo corticis Soymidæ ſcrupuli ex aquâ ter adhi-
“ biti alvum bis cierunt, morbumque levarunt. Cortex
“ repetitus ægro ſanitatem reſtituit.

“ Pridie Iduûm, Dec. A. D. 1791, L— miles Indicus,

† Vide p. 148.

quilled, and red barks, which the best judges imagine are from the same tree; the thick-
red

“ annos natus viginti tres, antecedente die, febre quotidianâ
“ affectus est. Cortice ter singulis intermissionibus adhibito,
“ alvus soluta est, morbusque mox remisit.

“ Pridie Iduûm Dec. S. N— miles Indicus, annos natus
“ quadraginta, iv. Non. Decemb. febre correptus erat. Nullis
“ hastenus medicamentis usus, magis nunc magisque debilis
“ evaserat. Cortex in remissione ter adhibitus ventrem
“ solvit, triduoque morbum depulsit.

“ Pridie Iduûm Dec. A. D. 1791. N— miles Indicus,
“ annum agens vigesimum quintum, pridie febre quotidianâ
“ affectus erat. Cortex Soymidæ, ter in unaquaque inter-
“ missione adhibitus, alvum movit, atque morbum brevî su-
“ peravit.

“ VIII. Cal. Martii, A. D. 1792. J. V— per biduum
“ febre iterum* laboraverat. Morbo autem duo accessus
“ totidemque remissiones quotidie erant, ejus instar paulò
“ suprâ descriptæ†. Cortex Soymidæ, in matutinis inter-
“ missionibus, alterâ quaque horâ adhibitus, triduo febrem
“ curavit.

“ Circiter medium Februarii, R— infector telæ xylinæ,
“ annos natus viginti quinque, laborans tumore hypogastrii
“ æquali, dolente, quem comitata est febris omni mane rece-
“ dens, atque alvus astricta, ad Roxburgiûm adductus est;
“ cui dixit, se duodecim antè dies affectum esse dolore circa
“ umbilicum torquente, qui uno alterove die gravis evasit,
“ atque profundus, et, quasi inter vesicæ urinarie fundum
“ atque intestinum rectum. sedem cepit; abdomen mox tu-
“ muisse, ipsumque, toto corpore febricitasse; causam autem

* Vide p. 145.

† Vide p. 147.

“ ignorasse

red fort being from the trunk, while the pale-
quilled fort is from the branches, and from
young

“ ignorâsse malorum; multa denique remedia vernacula
“ incassum adhibuisse.

“ Ei præcepit medicus, ut assumeret parvas lixivie tar-
“ tarisatæ quantitates, donec superveniret catharsis, pro
“ potu communi biberet aquam ex tamarindis coctam cum
“ saccharo, et ut interea diætâ ex oryzâ famem tolleret.

“ Alvo his exoneratâ, meliuscule se habere sensit æger;
“ tumori autem nequaquam decrescanti, vesicatorium ad-
“ motum est, alvusque lixivâ tartarisatâ et aquâ ex tama-
“ rindis cum saccharo commistâ soluta est.

“ Per noctem febris invaluit. Die autem, à curatione
“ inceptâ, tertio alvus vehementer fluxit. Dejectiones
“ purulentæ admodum erant, pessimè olentes, colore per-
“ virides. Tumor statim subsedit.

“ Æger maximè debilitatus, per noctem, graviter fe-
“ bricitabat. Mane igitur, cum primùm febris se remisif-
“ set, ei pulvis ex Soymidæ cortice et lixivâ tartarisatâ com-
“ positus adhibitus est, et, die progrediente, ter repetitus.
“ His factis, alvus purulenta quædam quater dejecit. Hâc
“ curatione triduo post à febre valebat, et, cortice nunc
“ semel tantum in die adhibito, decem diebus domum re-
“ diit sanus.

“ Roxburgius unam tantam occasionem corticis Soymidæ
“ contra gangrænam adhibendi nactus est. Viro dissoluto,
“ per idem tempus lue Venereâ laboranti, super mediam
“ tibiam ulcus erat. Cum Soymidæ pulvis eius stomacho
“ nigratus esset, extracto usus est, atque, expectatione ci-
“ tiùs, morbo immunis evasit. Perhibet præterea Roxbur-

young trees. The Spaniards themselves, he adds, employ the red fort.

Second.

“ gius, Duffinum chirurgum valetudinarii Madrasienfis primarium hunc corticem contra istiusmodi mala maximo cum fructu adhibuisse.

“ His memoratis, Roxburgius ingenuè fatetur insignem tempestatis siccitatem, hujus novæ Swietenix corticis usum feliciorē forsitan reddidisse. Notat præterea, corticem primo die alvum plerumque movisse, postea autem nunquam, neque profectò, præter morbi curationem, ullos ex ejus usu effectus observasse. Cur, ante corticis usum, non sæpius, ut mos plerisque est, vomitum et alvum movisset, hanc rationem reddit, nempe ex regionis naturâ, ex victu, ex vitâ, atque ex religione, corpora Indis esse gracilia, nec plena, nec inflammationibus obnoxia; atque remediis, quæ ante corticem adhiberi solent, febres, ut ille putat, in longum sæpe trahi, et iis, æque ac morbo fere ipso, ægros infirmari.

“ Hæc uberiùs dixi atque fusiùs cò quòd ex his potissimum, quantum polleat hic cortex, apparet. His adductus pater meus, cùm ægros nosocomio Edinburgensî curabat, atque discipulis de iis prælegebat, nova hujus corticis tentamina facere voluit. Hæc autem regione, cùm febris intermittens perrara sit, nobis nulla, quid proficiat cortex noster, experiendi idonea satis occasio oblata est. Nonnullis autem ægrotis adhibita est.

“ XIII. Cal. Jan. A. D. 1793. Joannes M'Kay, annum agens vigesimum tertium, priusquam in nosocomio receptus erat, duodecim dies febre, cujus accessiones alterò quoque die redibant, laboraverat. Sed, cùm, ab initio
“ horror

Second. *Cinchona Caribæa*; the Caribæan
or

“horror et calor per idem tempus duravissent, sudor prorsus defecisset, atque mala pectoris, coma, et torpor febrem comitata essent, hæc affectio minimè idonea, in quam novum medicamen tentaretur, videbatur. Cortex igitur Cinchonæ rubræ, per duodecim dies adhibitus est; cùm autem accessus post intervalla, licet valde dissimilia, adhuc redirent, ægro, ut Soymidæ drachmam alterâ quaque horâ fumeret, præscriptum est. Alvum torminibus magnopere movit, accessus autem proximus postremus erat. Convaluit.

“Jacobus Grant, annos natus viginti quinque, qui aliquandiu in nosocomio propter testis tumorem manserat, VIII Iduùm Junii, A. D. 1793, herbâ humidâ vespere recumbens, frigore, gravi dyspnœa atque angustia in faucibus sensu, affectus est. Hæc facile ætheri vitriolico cesserunt, cortexque Cinchonæ, quo vires prostratas reficeret, adhibitus est. v. Iduùm iterum frigore, dyspnœâ, atque vomitione sanguinolentâ, correptus est. Quinto postea vespere horrores, intermittentis instar, accesserunt. Usus corticis Cinchonæ, quippe qui accessionibus nihil obstaret, intermisit medicus, pulveremque corticis Soymidæ, duplici autem quantitate, in ejus locum adhibuit. Hoc factò morbus nunquam postea rediit.

“Duabus adolescentulis, alteri à singulari affectione hystericâ, convalescentibus cortex Swietenix Soymidæ, ut corpora firmaret, si non cum utilitate saltem sine incommodo, adhibitus est.

“Vi insuper astrictoriâ pollere, satis constat è muliere annorum quadraginta sex, quæ leucorrhœâ laborabat.

or Jamaica bark of Dr. Wright *. This last, our author observes, possesses in a higher degree the bitter, but is very weak in the astringent power, and ought not to be depended on when the other is procurable.

Third. *Cinchona Sanctæ Lucæ, floribus paniculatis, glabris, laciniis linearibus tubo longioribus, staminibus exertis, foliis ellipticis glabris*; Saint Lucia, or new bark. This is another sort, which has been introduced into practice: but its being possessed of strong emetic and purgative qualities, renders it, in our author's opinion, less eligible, particularly after the passages have been cleared. These properties, he observes, the Jamaica bark does not possess; which establishes a striking difference.

Fourth. *Cinchona Corymbifera, foliis oblon-*

“ Duobus senibus ventris fluxu affectis nihil profecit. Hi
“ autem, omnia, quæ alvum astringunt, experti, morbo
“ non levato, è nosocomio egressi sunt.

“ Cortex Soymidæ, ut multum, necne, contra putredinem posset, appareret, quinque ægrotis typho putrido
“ laborantibus adhibitus est. Omnes convaluere. His
“ ventrem adeo non movit, ut, per totum morbum, alvum
“ aliis medicamentis ducere opus esset.” *Vide Duncan Tentam. de Swietenia Soymida, p. 41. et seq.*—EDITOR.

* See Philof. Transact. Vol. LXVII. page 504; and London Medical Journal, Vol. VIII. page 239.

gis,

gis, lanceolatis, corymbis axillaribus; of Dr. Forster; is a native of the South-Sea Islands: but of its virtues we know nothing more, than that he says, “ it is like Peruvian bark, bitter “ and astringent.”

Fifth. *Cinchona Orixensis, foliis oppositis, tomentosis, stipulis interfoliaceis, semilanceolatis, floribus terminalibus, paniculatis, tomentosis, capsula valvis contrariis à vertice debiscens*; of Dr. Roxburgh. The structure of the capsule, he observes, forms the chief difference between this and *Cinchona Officinalis*, for the seeds are exactly as delineated by Gærtner, and the rest of the definition corresponds with that given by Linæus. It is a native of that chain of mountains which separates the northern provinces, or circars, from the Mahrattah dominions immediately behind them. The bark of this species likewise is bitter and astringent.

Dr. Roxburgh has also found another new species of *Swietenia*, a middle-sized tree, the wood of which is very heavy, close-grained, and yellow; the bark likewise is yellow, and very bitter, but possesses much less astringency than that of the *S. febrifuga*, and its astringency, he observes, is of a peculiar kind, for the colour produced, on an infusion, with a martial solution, was a dark brown.

There

There is also the bark of another large tree, which, at the time of writing this account, he tells us, he had under examination, and which is likewise very bitter: the Hindoos call it *Wallurse*. It will, he imagines, form a new genus in the class Decandria, and order Monogynia. Its essential characters are *calyx quinquefidus, petala quinque, nectarium duplex, exterius cylindricum oré decemfido, antheras gerens, interius annularium, basin germinis cingens, bacca monosperma*.

The bark of this tree, we are told, is in high repute as a medicine amongst the Hindoo physicians; and gives name to a compound soft extract, called *Walluvodusay*, which they employ in a variety of diseases.

It also possesses powers of a very different nature; for, powdered and thrown into pools where there are fish, it soon intoxicates them to that degree, that they are easily taken with the hand.

Dr. Roxburgh observes that the bark of *Melia Azadirachta*, already taken notice of*, has frequently and successfully been employed as a substitute for Peruvian bark, in the cure

* Page 133.

of remittents and intermittents; and that an infusion or decoction of its leaves is also a good anthelmintic, and as such employed by the Hindoos.

The bark of another large tree, which our author calls *Nauclea Daduga*, possesses also, he tells us, in a considerable degree, both the bitterness and astringency of Peruvian bark; and he thinks it is next in power to that of the *Swietenia febrifuga*. Although this tree differs widely in its flower from the hitherto known species of *Cinchona*, yet in its parts of fructification it agrees with them, it seems, almost exactly.

X. *An Account of the Effects of Mahogany Wood in Cases of Diarrhœa. By Mr. Francis Hughes, Surgeon of the General Infirmary at Stafford. Communicated in a Letter to Mr. John Pearson, Surgeon of the Lock Hospital, and by him to Dr. Simmons.*

AN accidental circumstance first suggested to me the idea that mahogany wood might prove serviceable as a medicine; for I did not then know that any part of the tree had been employed for medicinal purposes. I was accordingly induced to make use of it in cases of diarrhœa, both in decoction and in the form of an extract; and after repeated trials, I can venture to assert that I have not been disappointed in the expectations I had formed of its efficacy.

For the decoction I boil an ounce of the shavings of Jamaica mahogany wood in two pints of water, till one pint of the liquor is wasted, and then strain off the remainder for use.

The extract I make use of has been prepared
by

by boiling the shavings of the same wood in repeated affusions of fresh water, in the same proportion and manner as are directed for the extract of logwood (*extractum hamatoxyli*) of the London Pharmacopœia. The quantity of extract obtained in this way amounts to something more than $\frac{1}{8}$ of the shavings employed. The Honduras mahogany wood is of a paler colour, and less astringent than the Jamaica, and does not yield quite $\frac{1}{10}$ part of extract.

Both the decoction and extract are very bitter and astringent, leaving a roughness in the mouth for some time after they have been tasted.

The extract, in its appearance, resembles gum kino. It dissolves completely in water, and in spirit of wine, and strikes a black colour with salt of steel.

The following are some of the Cases in which I have employed these remedies.

CASE I.

In July, 1793, a soldier belonging to a regiment on the Irish establishment, who is a native of Stafford, was sent hither from his regiment

giment for the recovery of his health. He had for some time been unfit for duty, and was much reduced by a diarrhœa, which having come on after a fever, had continued several months, and resisted a variety of medicines.

I gave him an ounce of the decoction three times a day, and as it sat easy on his stomach, and seemed to have a good effect, the dose, after the third day, was increased to an ounce and a half. He persevered in the use of it during sixteen days; the diarrhœa gradually subsided; his appetite and strength returned; and at the end of that time he was sufficiently recovered to go back to his regiment in Ireland.

CASE II.

A woman of a thin, delicate habit, applied to me in October, 1793, on account of a violent diarrhœa, for which she had taken different medicines without any good effect. It had come on, she said, after sitting up a whole night in wet clothes, and had continued more than a fortnight; she was free from fever.

I directed her to take pills composed of six
grains.

grains of the extract, three times a day. Within the space of a week the diarrhœa was much abated, and she had acquired strength; she persevered, however, in the use of the medicine for the space of three weeks, at the end of which time the complaint had entirely ceased. A fluor albus, with which she had been troubled many months, was likewise much abated; but perhaps this latter circumstance ought rather to be ascribed to the improved state of her general health, than to any specific effect of the medicine.

CASE III.

In January, 1794, I was applied to by a man fifty years old, who for several years had been a hard drinker, and was now extremely emaciated; his legs were oedematous; he had no appetite; was subject to frequent vomiting, and had a slight diarrhœa.

I gave him aromatic bitters for several days, but finding no amendment, I determined to have recourse to the mahogany. I gave him eight grains of the extract, made into pills, three times a day. At the end of five days his disposition

disposition to vomit had ceased, and he had a little appetite. He continued the use of the medicine for ten days longer, and was then so much relieved as to be able to walk and ride out every day. This state of amendment continued for a fortnight, when he relapsed into his old habit of drinking, and his former symptoms returned. Recourse was again had to the same medicine, but without effect.

To the above I could add many other instances of the good effects of the extract and decoction in cases of long continued diarrhœa, where the complaint seemed to depend on a morbid irritability of the stomach and intestines, and where the use of tonic and astringent medicines appeared to be indicated. The few histories I have related will, I trust, be sufficient to point out the modes of administering the remedies in question, and the effects that may be expected from them; and perhaps will induce medical practitioners to extend a trial of their efficacy to other diseases.

The doses in which I have hitherto given these remedies have been small; but much larger doses may be given with safety, and in many cases will, I am persuaded, be more efficacious.

To

To try the effect of a considerable dose on the stomach, I took two ounces of a decoction, prepared by boiling two ounces of the shavings in two pints of water to a pint, which is twice the strength of the decoction described in Case I. and which I have usually administered. At first I perceived no effect from it; but at the end of ten minutes a disagreeable nausea came on, with a slight pain at the stomach, and a glowing sensation similar to that produced by the taking a glass of strong wine. These effects gradually went off in about half an hour, and I felt no other inconvenience from the dose.

Stafford, Feb. 12, 1794.

XI. *Account of some Discoveries made by Mr. Galvani, of Bologna; with Experiments and Observations on them. In two Letters* from Mr. Alexander Volta, F.R.S. Professor of Natural Philosophy in the University of Pavia, to Mr. Tiberius Cavallo, F.R.S.—From the Philosophical Transactions of the Royal Society of London, for the Year 1793. Part I. 4to. London, 1793.*

THE subject of the discoveries and researches, concerning which I am about to write to you, Sir, is *animal electricity*; a subject which cannot but be extremely interesting to you. I know not if you have yet seen the work of a Professor of Bologna, Mr. Galvani, which appeared about a year since, with this title; ALOYSII GALVANI *de Viribus Electricitatis in Motu Musculari Commentarius*. Bononiæ, 1791, in 58 pages, 4to, with four large plates; or at least if you have had any

* In the Philosophical Transactions these two letters are given in French; for the present translation of them we are indebted to the kindness of a friend.—EDITOR.

account of it*. It contains one of the most beautiful and surprising discoveries, and the germe of several others. Extracts from this work have appeared in different Italian Journals, and, among others, in that entitled *Gio-nale Fisico-medico*, published by Dr. Brugnattelli, of Pavia, to whom I myself have sent two long papers, which will be followed by several others, as I have considerably extended my experiments and inquiries on this subject. The letters I now address to you are intended as a sketch both of the admirable discovery of Mr. Galvani, and of the progress which I have been fortunate enough to make in this new path; and I request you, Sir, to present them to Sir Joseph Banks, Bart. the worthy President of the Royal Society, to be communicated, if he thinks proper, to that learned body, as a feeble testimony of my gratitude for the honour they have done me in electing me one of their number, and of my zeal and eagerness to comply with their invitation to communicate to them, from time to time, the fruit of my researches.

(1.) Mr. Galvani having dissected and prepared a frog, in such a manner that the legs remained attached to a part of the back bone,

* See Vol. III. p. 180.—EDITOR.

separated from the rest of the body, solely by the crural nerves, which were laid bare, observed that very lively motions were excited in these legs, with spasmodic contractions in all the muscles, every time that (this part of the animal being placed at a considerable distance from the conductor of an electrical machine, and under certain circumstances, which I shall explain hereafter) a spark was drawn from this conductor, not on the body of the animal, but on any other body, or in any other direction. The requisite circumstances, therefore, were, that the animal thus dissected should be in contact with, or very near some metal or other good conductor, of sufficient extent, or, what was still better, between two similar conductors, one of which should be turned towards the extremities of the legs of the animal, or some one of its muscles; the other towards the spine, or its nerves: it was likewise very advantageous that one of these conductors, which the author distinguishes by the names of *conductor of the nerves*, and *conductor of the muscles*, and preferably the latter, should have a free communication with the floor. It was in this situation especially, that the legs of the frog, prepared as above described, received violent shocks,

shocks, sprang up and contracted with vivacity at each spark drawn from the conductor of the machine, although it was at a considerable distance, and although the discharge was made neither on the conductor of the nerves, nor on that of the muscles, but on any other body, equally remote from them, and having any other communication through which the discharge might be transmitted, for instance, on a person placed in the opposite corner of the room.

(2.) This phenomenon surprized Mr. Galvani, perhaps more than it ought to have done; for the power, not only of electric sparks when they immediately strike the muscles or nerves of an animal, but of a current of this fluid traversing them, in any manner whatever, with sufficient rapidity, its great power, I say, of exciting commotions, was a thing sufficiently known; besides, it was obvious how, in this experiment, and in all those of the same kind, related in the first and second parts of his work, and which are represented in the two first plates of figures, his frog became liable to be affected by such a current. We have only to consider that well-known property of electrical atmospheres, or what is called *compressive electricity*, by which the fluid of conducting bodies,

placed within the sphere of action of an electrified body, is compressed and displaced, in proportion to the force and extent of this sphere, and kept in this state of displacement so long as the electricity subsists in the predominant body; and when this is removed, returns to its place gradually, if the electricity of that body is slowly dissipated, or in an instant if it be destroyed instantaneously, by discharging suddenly the body that contained it. It is this returning current, therefore, this reflux of electrical fluid in the conducting bodies contiguous to the frog, or near it, its sudden passage from the conductor of the muscles to the conductor of the nerves, or *vice versa*, through its body, especially when such a current is compressed in the single and narrow channel of the nerves, which excites the spasms and movements in the experiments in question. Mr. Galvani, who seems not to have sufficiently reflected on this property of electrical atmospheres, and who was not aware of the prodigious sensibility of his frog, singularly prepared in the manner above described (I must here observe that I have found this sensibility nearly equal in all the other small animals, such as lizards, salamanders, and mice) was extremely struck with such an effect, which will probably

probably not appear so marvellous to other philosophers. This, however, was the first step which led him to the grand and beautiful discovery of an animal electricity, properly so called, and which belongs not only to frogs and other animals of cold blood, but likewise to every animal of warm blood, quadrupeds, birds, &c.; a discovery which forms the subject of the third part of his book, a subject altogether new, and very interesting. It is thus he has opened to us an immense field, into which I propose to enter, and pursue my researches, after I shall have dwelt a little more on those preliminary experiments which relate to the action of artificial or extraneous electricity on the nervous and muscular fibres.

(3.) It was chance that presented to Mr. Galvani the phenomenon I have been describing, and which astonished him (I repeat it) more than it ought to have done. Still who would have believed that a stream of electricity, so feeble as not to be rendered sensible by the most delicate electrometer, should be capable of affecting so powerfully the organs of an animal, and of exciting in its limbs, cut off one or more hours before, movements, nowise inferior in strength to those produced in the living animal,

animal, such as vigorously darting out its legs, springing up, &c. to say nothing of the most violent tonic convulsions? And yet such is the stream that affects the little animal, placed, for instance, on a table, near some metal, or between two good conductors, not insulated, when a person draws from the prime conductor, suspended several feet above, a moderate spark, and conveys the discharge through quite another channel.

(4.) I say *moderate*; for if it be very strong, and the conductor, large and highly charged, be not at a very considerable distance from the bodies on the table, little sparks will be perceptible in the interstices of these bodies, especially the metallic ones, and even in the place where the frog forms a ring of communication between them, which sparks are evidently produced by the returning stream of electricity, of which I have already spoken, (sect. 2.) Or if matters do not come to this point, instead of sparks we may perceive movements, sufficiently obvious, of electrometers placed on the same table and in the same places. In this case, therefore, where the electrometer affords the sign, and much more in the other, where the above-mentioned sparks are obtained, we may observe,
that

that even a frog, entire and untouched, or any other small animal, as a lizard, a mouse, or a sparrow, is seized with strong convulsions in all its limbs, especially in its legs, which dart forwards with vivacity, if the passage of the electric fluid (the returning stream) follows the direction of these same legs from one end to the other. So far there is nothing wonderful; the circumstance that may excite surprise is in the case where the stream of electricity, though no longer sensible, not even to the most delicate electrometer, continues to excite the same convulsions, the same movements, if not in the entire frog, at least in its limbs, when dissected and prepared in the manner practised by Mr. Galvani.

(5.) I have endeavoured, with much attention, to determine what might be the least electrical power requisite to produce these effects, as well in the entire and living frog, as in one dissected and prepared in the manner above described, which is what Mr. Galvani has omitted to do. I have preferred the frog to every other animal, because it is endowed with a very durable vitality, and it is very easy to prepare it. I have, however, made experiments on other small animals with the same view, and with a success nearly similar. In order to estimate well
the

the strength of the stream of electricity, I have thought it right to submit the animal intended for experiments of this kind, not to the returning streams occasioned by electrical atmospheres (Sect. 2.), but to direct electrical discharges, sometimes from a simple conductor, sometimes from a Leyden phial, and in such a manner that the whole stream must have passed through the body of the animal. For this purpose I was careful to keep it insulated in one way or other, and most frequently by fixing it, with pins, to two flat pieces of soft wood, supported by glass columns.

(6.) I have found then, that for the living and entire frog the electricity of a simple conductor, of a middling size, is sufficient, when it comes only to be able to give a very weak spark, and to raise Henley's electrometer from five to six degrees; that if I make use of a Leyden phial, likewise of a middling size, a much weaker charge of this produces the effect, such a one, for example, as yielding not the least spark, and being nowise sensible to the quadrant-electrometer, is scarcely sufficiently so to Cavallo's electrometer to separate its little pendulum about 1-tenth of an inch.

(7.) This, as I have just now shown, for a
 2 frog

frog entire and untouched ; for when it is dissected and prepared in different ways, and particularly after Galvani's manner, in which the legs are connected with the dorsal spine merely by the crural nerves, a much weaker degree of electricity, whether from the conductor or from the Leyden phial (the fluid being obliged to pass through the narrow passage of the nerves), fails not to excite convulsions, &c. Yes, an electricity forty or fifty times weaker, as a charge of the phial that is absolutely imperceptible to the last-mentioned electrometer (Cavallo's), and even to that extremely delicate one of Bennet ; a charge, that I was able to render sensible only by means of my condenser, and which I think may be estimated at five or six hundredths of a degree of Cavallo's electrometer.

(8.) Thus then, in the legs of a frog attached to the spine of the back solely by its nerves (these being laid bare), we have a new species of electrometer ; since electrical charges, which from their yielding no sign to the electrometers already in use, would seem null, afford such obvious ones to this *animal electrometer*, if I may be allowed the expression.

(9.) When

(9.) When we have seen how, in a frog thus prepared, strong convulsions are excited by an extremely weak electricity, by an imperceptible stream of fluid, we ought surely to be no longer surpris'd, that the animal should be affected in the same manner when any body whatever discharges suddenly the prime conductor of an electrical machine, and occasions another stream of electric fluid, great or small, of the fluid before displaced in the conducting bodies near the frog, and which re-establishes itself, in the manner already explained (Sect. 2.), to pass rapidly through its nerves. Let us suppose this returning stream to be scarcely equal to that which a conductor, sufficiently bulky, throws off directly, with an electricity that yields no spark, and that is almost insensible even to Cavallo's electrometer, or a small Leyden phial, charged scarcely a tenth of a degree of this same electrometer; let us suppose, I say, that the stream of electricity is not stronger than this, still it will be sufficient, as my experiments, above related (Sect. 6. and 7.), show, to excite the movements in question.

(10.) But if, after the experiments just now referred

referred to, we ought no longer to be surpris'd at those of Mr. Galvani, described in the first and second parts of his work, how can we avoid being so at those entirely new and wonderful ones related in the third? Experiments in which he obtained the same convulsions and violent movements of the limbs, without having recourse to any artificial electricity, or extraneous excitement, by the simple application of a conductor, one end of which was made to touch the muscles, and the other the nerves or spine of the frog prepared in the manner already described. This conductor, he found, might be either entirely metallic, or composed partly of metal and partly of other bodies of the class of conductors, as water, one or more persons, &c. Even wood, the walls and floor of the room, might enter into the circle provided they were not too dry; it was only by the interposition of non-conducting substances, as glass, rosin, and silk, that the effect was prevented. Bad conductors, however, did not do so well, and only during the first moments after the animal was prepared, and so long as the vital powers remained in full vigour; after which good conductors only were found to succeed,

ceed, and in a short time it was found impossible to produce the effect unless with excellent conductors, that is, with conductors entirely metallic. He moreover found a great advantage from applying a sort of metallic armour, or coating, to that portion of the spine which he left attached to the crural nerves, and to the nerves themselves, and particularly from covering this part with a thin leaf of tin or lead.

(11.) Mr. Galvani did not confine himself, in these truly astonishing experiments, to frogs; he extended his trials with success, not only to several other animals of cold blood, but likewise to quadrupeds and birds; in all of which he obtained the same results, by means of the same preparations, which consisted in laying bare some principal nerve at the part where it passes into a limb susceptible of motion, and after arming the nerve with some metallic substance, forming a communication, by means of his conductor, between this coating and the muscles to which the nerve is distributed.

(12.) It was thus he fortunately discovered, and demonstrated to us, in the most evident manner, the existence of a real *animal electricity* in all, or almost all animals. It seems in fact to be proved by his experiments, that the electric fluid

fluid tends incessantly to pass from one part to another of a living organized body, and even of detached limbs, so long as any remains of vitality subsist in them; that it tends to pass from the nerves to the muscles, or *vice versa*, and that the muscular movements are owing to a similar transfusion, more or less rapid. In truth, it would seem that no objections can be raised to this, or to the manner in which Mr. Galvani explains it, by a kind of discharge similar to that of the Leyden phial. But a great number of new experiments that I have made on this subject, will serve to show that many restrictions must be made with regard both to the thing itself, and to the deductions the author has drawn from it; my experiments likewise will be found considerably to extend the phenomena attributed to this animal electricity, and will display it to us under a great number of new circumstances and combinations.

(13.) Mr. Galvani, pursuing the idea he has formed to himself from his experiments, and adhering in every respect to the supposed analogy of the Leyden phial, and his conductor, imagines there is naturally an excess of electric fluid in the nerve, or in the interior part of the muscle, and a corresponding defect of this
fluid

fluid in the outer part, and *vice versa*; and he supposes consequently that one end of this conductor must communicate with the nerve, which he considers as the conducting wire or hook of the phial; and the other end with the external surface of the muscle. All the figures of his third and fourth plates, and all his explanations relate to this. But if he had a little varied his experiments, as I have done, he would have seen that this double contact of nerve and of muscle, this circuit which he imagines, is not always necessary. He would have found, as I have, that the same convulsions, the same movements may be excited in the legs and other limbs of frogs, and of every other animal, by placing metallic substances in contact with two parts of a nerve only, or with two muscles, or even with different parts of a single and simple muscle.

(14.) It is true we are very far from succeeding so well in this way as in the other, and that in this case it is necessary to have recourse to an artifice, of which we shall have occasion to speak more fully hereafter, and which consists in employing two different metals; an artifice, which is not absolutely necessary when the experiment is conducted according to Galvani's
method

method above described (Sect. 10 and 11), at least so long as vitality remains in full vigour in the animal, or in its detached limbs; but, at any rate, since by arming the nerves only, or the muscles only, with different metals, we are able to excite contractions in the latter, and movements in the limbs, we must conclude that if there are cases (and this may perhaps still be very doubtful) where the pretended discharge between nerve and muscle (Sect. 12 and 13.) is the cause of the muscular movements, there are likewise many and more frequent circumstances, where the same movements are obtained by quite another play, quite another circulation of the electric fluid.

(15.) Yes, it is quite another play of the electric fluid, of which we may be said rather to disturb than to restore the equilibrium, inso-much as it passes from one part to another of a nerve, a muscle, &c. as well internally by their conducting fibres, as externally by means of the metallic conductors that are applied, not in consequence of any respective excess or defect, but by a peculiar action of these same metals, when they are of different kinds. It is thus I have discovered a new law, which is not so much a law of animal electricity as a law of

common electricity ; to which we must attribute the most part of the phenomena, which, from the experiments of Galvani, and from several others which I made myself, seemed to belong to a true spontaneous animal electricity, but which in truth do not : they are really the effects of a very feeble artificial electricity, which is excited in a way never before suspected, by the simple application of two coatings of different metals, as I have already hinted, and which I shall explain better elsewhere.

(16.) I think it right here to say, that at the discovery of this new law, of this, till now, unknown artificial electricity, I was mistrustful of every thing that seemed to me to demonstrate a natural electricity, in the strict sense of the term, and that I was on the point of giving up this idea. But upon carefully reconsidering all the phenomena, and repeating the experiments under this new point of view, I found that some of them support such an idea, (those, for instance, in which there is no need of different coatings, or even of any coating, a simple metallic wire, or any other conducting body, performing the office of conductor between the nerve, and one of the muscles connected with it, being capable of exciting convulsions in the latter).

latter), (Sect. 10, &c.) and that thus a natural animal and properly organic electricity subsists, and cannot be entirely overturned. The phenomena which establish it, although much more limited, are however sufficient to demonstrate its existence, as I have just now mentioned, and as will more clearly be shown hereafter.

(17.) What will perhaps be found more disagreeable is, that we must likewise confine within narrower limits its influence in the animal œconomy, and give up the finest ideas we had formed of it, and which seemed to be about leading us clearly to explain muscular motion. My experiments, varied in every manner possible, show that the motion of the electric fluid excited in organs, does not act immediately on the muscles; that it does nothing more than excite the nerves, and that the latter, put into action, excite in their turn the muscles. What this action of the nerves is; how it propagates itself from one part of a nerve to another; how it passes to the muscles, and how the motion of the latter results from it; these are problems, in the explanation of which we are not farther advanced than before the discovery in question.

(18.) I come now to the experiments that prove all the assertions I have advanced in these

last paragraphs. From a great number I shall select only a few, which seem to me the best calculated to establish certain principles, for the most part new and different from those adopted by Mr. Galvani. But I must first say a few words more concerning the experiments of this writer. I know not whether he has made others, but those he has described in his work are included in too narrow a circle; in all of them the object is to lay bare and insulate the nerves, and to establish a communication, by means of conducting bodies, between these nerves and the muscles that are dependent on them (as may be seen in all the figures of the four plates annexed to his work), in order to excite convulsions and movements of the muscles, by the action of the electric fluid. He supposes therefore, in every case, and he explains himself pretty clearly on this point, that the transfusion of the electric fluid that is produced, whether by artificial electricity, or by natural animal electricity, must take place from the nerves to the muscles, or *vice versa*; that these two limits at least must be included in order for the muscular movements to take place; and in truth all the experiments he has described seem to prove this. But then they are confined, as I have just now said,

said, within a circle that is too limited, and beyond which he has never, or scarcely ever, extended his inquiries. By varying the experiments of this kind in different ways, I have shown, that neither the one nor the other of those conditions, viz. the laying bare and insulating the nerves, and the touching simultaneously these and the muscles, in order to procure the supposed discharge, are absolutely necessary (Sect. 13.). It is sufficient, when, for instance, we have laid bare the ischiatic nerve of a dog, lamb, &c. if we pass a stream of electricity from one part of this nerve to the other, even though it be near, and leave all the rest untouched and free; it is sufficient, I say, to do this in order to excite in the limb very strong convulsions and movements; and this whether we employ an extraneous artificial electricity, or excite the electric fluid that is inherent in the nerve itself. Here is the manner in which I make these experiments.

(19.) EXPERIMENT A. I compress, with a pair of forceps, the ischiatic nerve a little above its insertion into the thigh, and I apply, a few lines higher up, a piece of money, or a plate of metal, on this same nerve, carefully separated from the parts that adhere to it, and supported

by a thread, a plate of glass, a stick of sealing wax, a piece of dry wood, or any other substance that is a bad conductor. Then placing the belly of a Leyden phial, very weakly charged, on the forceps, I bring the hook into contact with the other piece of metal; and the moment the discharge takes place, although it be too feeble to produce the least spark, convulsions take place in all the muscles of the thigh and leg, the whole limb being agitated and springing up with more or less violence. And yet the whole of this leg, and even a part of the nerve which passes to it, are, as we see, out of the track which the electric fluid takes in its passage, so that only a small portion of the nerve can have been irritated; and yet this is sufficient to occasion the convulsion of the muscles.

(20.) EXPERIMENT B. The same effects, that is to say, similar convulsions and motions of the leg, take place, without our having recourse to an extraneous electricity, by the discharge which takes place, in a certain manner naturally, when we apply, as above, the same forceps, or a plate of silver, to one part of the nerve, and a plate of some other metal, and above all, of tin or lead, to another part, and then bring about a simple communication between

tween them, either by an immediate contact, or by the interposition of a third piece of metal made to perform the office of a conductor.

(21.) Thus we see that the same effects, that is, convulsions and violent muscular contractions, take place without any discharge of electric fluid between the nerves and muscles, in the manner Mr. Galvani supposes; and without requiring one end of a conductor to communicate with the one, and the other end with the other. Neither is the other condition, that of laying bare the nerve, and freeing it of its adhesions, at all more necessary, as will appear from the following experiments.

EXPERIMENT C. I apply coatings, or plates, of different metals; (and it is this difference of coatings that is essential) (Sect. 14. and 15.) to an entire and living frog, that is covered with its skin, and, in short, is untouched. I apply, for example, a thin piece of tin-foil on its back, or its loins, and I place a piece of silver money under its thighs, or its belly, slightly compressing it; this done, I slide forward the piece of money till it comes into contact with the tinfoil, or I form a communication between the two metals by means of a piece of iron wire, or any other metal; and at that instant convulsive motions

N 4

take

take place in all the muscles of the belly, thighs, and back, with violent tremors of the legs, contraction and curvature of the spine, &c. which convulsions and spasms, although nearly universal, are however most considerable in the limbs and muscles contiguous to the coatings, and still more so in those which are dependent on the principal nerves nearest to the two metals.

(22.) These experiments succeed in some other animals; in fishes, and particularly in eels, in none of which is it necessary to remove the skin, though it does not fail, in a small degree, to lessen the effect. This is why, by removing it, at least in part, particularly in the frog, we obtain the effects with more certainty, and to a greater degree. We likewise gain something, in this respect, by cutting off the head of the frog, and thrusting a large pin into the spinal marrow; we then excite, by means of different coatings in the manner above described, stronger movements, or at least such as are more obvious, because they are no longer confounded with the movements the animal gives itself while living.

(23.) If it be advantageous, as we have seen, to take off the skin of frogs, although very thin
and

and pretty moist, it is much more so, and even necessary, to remove it from almost all the other animals, as lizards, salamanders, serpents, tortoises, and more especially from quadrupeds and birds, that are furnished with a drier and much thicker skin, to succeed in these experiments. The following, therefore, is the mode I adopt.

EXPERIMENT D. I fasten to a table, by means of some large pins, a lizard, a mouse, a fowl, &c. and after making an incision through the skin, and other integuments, to the bare flesh, on the back of the animal, I turn back the integuments on each side; I do the same on the thigh or the leg; after which I apply the two metallic coatings on the exposed parts, viz. on one the tin foil, and on the other a spoon or a piece of money; I then form a communication between the two coatings, and every time I do this I excite strong contractions in the adjacent muscles, and particularly in those of the thigh and leg, which moves and agitates itself with great violence. These convulsions are much more considerable when the tin foil is applied near the ischiatic nerve, and the piece of silver on the gluteus muscle, or on that named gastrocnemius; and the effects are still greater if the nerve itself is laid bare, and coated with
the

the tin foil; if, leaving it attached only to the muscles to which it is distributed, we deprive it of every other adherent part; or if, in short, we separate the entire limb from the rest of the body, with its nerve hanging out, and submit it in this state to our experiments.

I am, &c.

A. VOLTA.

September 13, 1792.

SECOND LETTER.

(24.) It will be sufficiently understood that what I have said with respect to the ischiatic nerve, and the leg, is applicable to the brachial nerve and to the arm, as well as to every other nerve relatively to the muscles under the influence of that nerve.

(25.) These last preparations are analogous to those of Mr. Galvani; and they clearly prove that it is advantageous to lay bare the nerves, and still more so, to detach them all round from the adherent parts; but they are far from showing

ing that this is a necessary condition, since we never fail to obtain the same convulsions and movements of the limbs when we simply lay bare the muscles, and leave the nerves covered and concealed under them in their natural state, as all my other experiments above related (Sect. 21, 22, 23.) serve to show.

(26.) After these trials on reptiles, birds, and small quadrupeds, I proceeded to other and larger animals, as rabbits, dogs, lambs, and bullocks; and I not only succeeded in obtaining similar effects in all the ways above described, but even stronger and more durable ones, by reason that the vital heat maintained itself in those large animals, and in their limbs, a longer time. For I ought not to omit to say, that if in the most part of animals of cold blood, and particularly in frogs, the vital principle subsists in detached limbs several hours, that principle which renders them so sensible to the weakest electrical irritation, it hardly continues beyond a few minutes in animals of warm blood, and commonly disappears before the whole of this animal heat is dissipated.

(27.) Having had such success with my experiments on large and small animals of every kind, in some instances alive and entire; in
others

others deprived of their skin, or their head, or dissected in different ways ; and having obtained similar effects in their large detached limbs, and almost always without the preparation required by Mr. Galvani, that is to say, without laying bare the nerves, I was desirous of going still farther, and of making similar trials on smaller limbs, on a single muscle, and even on small portions of muscles ; and the fresh success I had in these trials led me to other discoveries, which I will soon mention, after having described some of these experiments.

(28.) EXPERIMENT E. I cut off, in some instances, the leg and thigh of a frog, in others, the leg only, and in some half or a quarter of a leg ; and on applying, as usual, to one part of the amputated portion the tin foil, and to the other the plate of silver, and forming a communication between these two coatings, I constantly excited convulsions and movements ; I have even separated a single muscle, for instance the gluteus, or the gastrocnemius, and sometimes only a portion of muscle not larger than a barley-corn, and yet the same effects, that is to say, very strong contractions of these muscles, or parts of muscles, have been produced by means of two different coatings, &c.

EXPE-

EXPERIMENT F. I have repeated the same experiments on a leg, on a half or a third part of the leg, on a single muscle, or part of a muscle, of a fowl and other birds; on a slice of the gluteus of a rabbit, a lamb, &c. and I have had the same effects as long as the flesh preserved a sensible heat. (Sect. 26.)

(29.) Thus then we are able to excite very strong contractions in the muscles of animals of warm as well as of cold blood, and in every detached portion of muscular flesh; and this by means of the simple artifice of different metallic armours or coatings, applied to the muscle itself, without any preparation of the nerves, and even without laying them bare. We have besides seen that we can excite these effects quite as well, and by the same means of metallic coatings applied to two neighbouring parts of the same nerve, (Sect. 19, and 20. Experiments A. and B.) whence I have reason to conclude, that there is no necessity for a discharge of electric fluid to take place between nerve and muscle, or for any transmission of it from the interior to the exterior part of the latter by means of the nerve and metallic conductor, as Mr. Galvani supposes, or *vice versa*: and that there is no comparison to be made between

tween the muscle and the Leyden phial and its discharge, in the experiments in question. In fact, what resemblance or analogy is there to the Leyden phial, where the two plates of metal, a communication between which is formed by the conductor, are applied very near to each other on the external surface of the same nerve, (Experiments A. and B.) or on the external surface of two muscles, or even of the same muscle (Experiments C. D. E. F.); it must be confessed it would be in vain to attempt to support any analogy between any of these experiments and the Leyden phial.

(30.) EXPERIMENT G. Having placed two coatings, one of silver leaf, the other of tin foil, on exactly corresponding parts of the two thighs of a frog, I excited contractions of the muscles, and the usual motions of the legs, at the instant I formed a communication between the two coatings by means of the conductor.

(31.) Is it thus, I ask, that the discharge of two Leyden phials takes place, by forming a communication between their homologous surfaces? Let us lay aside, therefore, these ideas of phial and discharge, and every forced explanation, and let us say simply, that in these and other analogous experiments, a transmission of

the electric fluid takes place from one to another of two parts properly coated; a transmission determined, not by a *relative excess* of this fluid, which cannot naturally be supposed between parts that are similar, but by the diversity of these same coatings, which must be of different metals, as I have taken care already to point out, (Sect. 20, and 21. Experiments B. and C.) and uniformly to inculcate in the subsequent parts of my paper.

In fact,

(32.) EXPERIMENT H. If two muscles, or two parts of the same muscle, are similarly coated, that is, with two plates of the same metal, both of them equal in temper and hardness, in softness or rigidity, in the roughness or smoothness of their surface, and both are applied in the same manner, it will be to no purpose to bring about a communication between them by means of a conductor, as no convulsion, no motion will take place.

(33.) I confess it is not easy to conceive how and why the simple application of two dissimilar coatings, I mean of two different metals, to similar parts of the animal, and even to two parts very near to each other of any one muscle, shall disturb the equilibrium of the electric fluid,

fluid, and drawing it from its state of repose and inactivity, shall induce it to pass incessantly from one part to another ; which transflux takes place as soon as a communication, by means of the conductor, is formed between these two dissimilar coatings, and continues all the time this communication subsists. But conceivable or not, and whatever may be the cause, it is a fact that the experiments I have already related sufficiently prove, and which will be confirmed by many others, to the description of which I shall endeavour to add some explanation. It is a fact, to be added to what we already know in electricity ; a fact which must surely appear extraordinary, and difficult to be reconciled with the laws commonly established. It is truly a new and very singular law, which I have discovered ; a law that belongs not properly to animal electricity, but to common electricity, since this transflux of the electric fluid, a transflux, not momentary, as a discharge would be, but which continues as long as the communication between the two coatings subsists, and takes place whether these coatings are applied to living or dead animal substances, or to other conductors not metallic, but sufficiently good, as water, or moist bodies. But before I proceed to the experiments

periments which decisively prove all that I have advanced, I think it right to offer a few more remarks on those I have already described (Sect. 20—32.).

(34.) It would seem from these that by means of the simple artifice of coatings of different metals suitably applied, we are able to excite very strong convulsions in every muscle of every animal, so long as it continues to possess any degree of vitality. Such a conclusion, however, would be too general, my experiments having taught me that it is to be admitted only with certain restrictions, as well with respect to the classes and genera of animals, as with respect to the different muscles of each animal.

(35.) And first with respect to the different classes of animals; although it has uniformly happened that all the quadrupeds, birds, fishes, reptiles, and amphibious animals, which have been submitted to my experiments, exhibited the phenomena above described, it is no less certain that worms in general, and several species of insects, remained unaffected. I have in vain tried with worms, leeches, snails, oysters, and different caterpillars; I have not even been able to excite the least motion in them by small and moderate sparks, and discharges of artificial

electricity. Here is the manner in which I proceeded.

EXPERIMENT I. I applied the tin foil, and silver leaf, to different parts, as well external as internal, of these snails, leeches, earth worms, &c. and in the best way I was able; I then formed a communication between these metallic coatings, sometimes by bringing them into contact with each other, and at others by means of another metal that performed the office of a conductor; but by neither of these means could I ever obtain the least motion in any part of the body.

EXPERIMENT L. I conveyed through their bodies, both when insulated and not insulated, discharges of a Leyden phial of sufficient strength to excite a moderate spark, and to give me a slight shock, but they were not sensibly affected by it; no motions or convulsions were produced.

(36.) Does it follow from hence that the more imperfect animals, the whole class of worms, and several species of insects, are destitute of that sensibility and irritability, that electrical mobility, if I may be allowed the expression, with which other more perfect animals are endowed? I am unwilling to draw this general conclusion from my experiments, because I have

as yet extended them only to a small number of worms and insects; and with regard to the latter, I think it right to observe that I have succeeded, without much difficulty, with crawfish, beetles, grasshoppers, butterflies, and flies. It may not be useless that I explain one of the ways in which I succeed with these animals, as they are with difficulty submitted to experiments, on account of their minuteness, or of the scales with which they are covered.

EXPERIMENT M. After cutting off the head of a fly, a butterfly, beetle, &c. I slit open, with a penknife or small scissors, the whole length of the corselet, and introduce deep into the slit, near the neck, a bit of tin foil, (what is improperly called silver paper is very fit for this purpose) and a little below I introduce, and likewise deep into the slit, a bit of silver plate, or small silver coin; and when I bring the latter into contact with the piece of tin foil, the legs begin to bend and tremble, and the other parts, and even the trunk of the animal, are thrown into agitation. It is very amusing to excite in this manner the chirping of a grasshopper, &c.

(37.) After what I have just now said, I should be wrong to rank insects among the animals that are destitute (like the class of worms

above mentioned) of the electrical property in question. At the utmost, if caterpillars appear to be so, it may be said that in this state of larva, before they have attained, by their metamorphosis, a perfect state, and acquired new organs, &c. they may be compared in many respects to worms, and, like these, are not endowed with electric sensibility.

(38.) In short, if I may be allowed to state here what I think, those animals only that have very distinct limbs, with joints, and muscles fitted for the motion of those joints, or, in other words, muscles that are called flexors, or levators, and nerves proper to regulate them, such animals only, I say, are sensible to, and become seized with real spasmodic contractions in consequence of either small discharges of artificial electricity, or a weak current of fluid occasioned simply by different metallic coatings; which contractions and spasms bring on the motion, and even a violent agitation of the said limbs. On the contrary, worms, and such insects as have not sufficiently distinct limbs, or joints properly so called, or which are destitute of flexor muscles, or enjoy only a vermicular motion, are nowise affected by such an electricity. The motions of these animals depend on
a different

a different animal œconomy; on a different mechanism, which in several species has been very well discovered and explained. Such are my ideas, still indeed somewhat vague, and founded only on a few experiments; it is the sequel of these that must either confirm or rectify them.

(39.) With respect to different muscles in the same animal, I am able to advance something more certain. I say then, that all muscles are very far from being susceptible of contraction from the weak electricity in question. There is a great distinction to be made with regard to their functions in the animal œconomy; all of them are not subject to the empire of the will, and fitted for spontaneous movements: and, strictly speaking, it is only those which are so that are capable of spasmodic contractions by the means above described; yes, the muscles subject to the will are the only ones I have found susceptible of irritation and motion, by the action of that weak current of electric fluid occasioned by the simple contact of two different metals. The other muscles, over which the will has no direct power, as those of the stomach, intestines, &c. are not at all so, not even the heart, though in other respects so irritable. We must except, however, the muscles of the

O 3

diaphragm,

diaphragm, (and I conjectured it before I made the trial) these being of the number of those whose motion depends on the will.

EXPERIMENT N. It is very surprising that a slice of good muscular flesh, cut, for instance, from the thigh of a lamb killed half an hour or an hour before; that this piece, I say, of muscle, almost quite cold, and which is no longer sensible to the action of any mechanical or chemical stimulus, should be so powerfully affected by the electric fluid conveyed from one part of it to another, as to be seized with very strong spasmodic contractions; and that, on the contrary, the heart recently taken out of the same animal, and still warm and very irritable, should, when treated in the same manner, with the best adapted metallic coatings, suffer no alteration upon our making a communication between the two metals by means of the conductor; and that its pulsations, when weakened or slackened, or altogether suspended, should not be increased, or even revived, notwithstanding all this takes place from the application of the slightest mechanical or chemical stimulus.

(40.) The electric fluid, therefore, which seems to be the stimulus appropriated to the muscles of the will, is nowise so to the heart,

or

or to the other muscles formed for involuntary vital and animal functions. But what will be said if I make it appear that it is not the immediate or efficient cause of motion in the voluntary muscles; that even in these it is a mediate cause only, the nerves alone being directly affected by it? And yet this is what I have learned from several experiments; experiments that have obliged me to give up the finest and most extensive ideas I had formed on the subject. I was fond of thinking, with Mr. Galvani, that as often as a current of the electric fluid, put in motion in the organs, was impelled with a certain degree of strength to the muscles, this fluid did itself perform the office of a stimulant, and excited the irritability which is peculiar to them; that every muscular movement was executed in consequence of a similar irruption of electrical fluid into the muscles, either by means of artificial electricity, or by putting in motion the natural artificial electricity; that, in short, even the motions which are performed naturally in the living animal machine, at least the voluntary motions, acknowledged the same cause, that is to say, the immediate action of the electric fluid on the muscles. But I repeat it, I have found myself obliged, with regret, to

give up all those fine ideas by which it seemed possible to explain things to admiration. Yes, we must considerably limit the action of electricity in animals, and consider it under another point of view, that is to say, as being capable of exciting, of itself, the nerves, as I have already hinted, and as I shall now proceed to prove.

(41.) In the first place, then, that it can act, and that it really does act, on the nerves, and that the latter, excited by it, excite in their turn the muscles connected with them, without even the electrical stream's arriving at those muscles, is a fact which no longer stands in need of proofs after those furnished by the experiments A. and B. (Sect. 19. and 20.) and even by an experiment of Mr. Galvani, which, according to his account, was the first he made, and the origin of all his other experiments. It is sufficiently obvious that the electric current, in the experiment in question, as well as in those made by me, and which I have just now referred to, pervades only a part of the crural nerve, but not one of the muscles of the leg; and yet as the latter depend on the nerve, they are affected with convulsions.

(42.) But I go farther, and maintain, that
even

even in the cases where the electrical current (it will be clearly understood that I am speaking only of weak artificial discharges, or of the current which takes place by the simple application of coatings of different metals) strikes and penetrates muscles susceptible of movement, it is not by irritating the latter immediately that it occasions them to contract, but by stimulating their nerves. This is what is shown by my experiments C. and D. (Sect. 21. and 23.) where, upon the tin foil and piece of silver being applied immediately to the muscular parts of the animal, whether the animal or only a detached portion of it is the subject of the experiment, it is not so much the muscles covered by the two metallic coatings that suffer the most violent contractions, as those which depend on some principal nerve, to which one or other of the coatings is contiguous. It is in this manner that in the frog, when the tin foil is applied on the loins, where the crural nerves lay at but little depth, the muscles of the legs are seized more than any others with strong convulsions, more so even than those contiguous to the other coating, that is to say, to the piece of silver. I have already pointed out the same thing in quadrupeds, dogs, lambs, &c. with regard

gard to the ischiatic nerve, (Experiment D.) and I have only to add, that the leg never fails to be convulsed when this nerve does not lay too deep under the flesh and other integuments, and one of the coatings is properly applied to this part ; even although the other coating should be made to correspond neither with the gluteus nor any muscle of the leg, but with any other muscle whatever, provided it be not at too great a distance. Here is another proof why this happens :

EXPERIMENT O. If we apply in a frog, or any other small animal, the tin foil the whole length of the spine of the back, from which proceed all the nerves of the trunk and limbs, and the other coating to any other part whatever, all the limbs become affected ; the muscles, not only of the legs, but of the belly and back, experience spasmodic contractions, and the trunk itself becomes curved ; in a word, the convulsions are general. The experiment is still more striking in a lizard than in a frog, and I shall therefore describe it.

EXPERIMENT P. After cutting off the head of a lizard, and laying bare the muscles of the back by removing the skin, I apply a piece of tin foil to the mutilated end, in such a manner
that

that the tin foil is spread beyond the edges of the wound, so as to rise a little over the shoulders, and I place a piece of money on the middle of the spine; this done, I slide forward the piece of money till I bring it into contact with the tin foil. At that instant the legs move, the tail twists itself, and the whole body of the animal becomes agitated, and darts from right to left, and from left to right. Is not this because the upper part of the spinal marrow, the principal source of the nerves, is irritated?

(43.) Nearly the same effects may be obtained by a similar operation on a mouse, a small bird, &c. but in these it is necessary to remove not only the skin and other integuments, but likewise some of the flesh, and this because their back being more fleshy, the principal nerves of the spine are more concealed by this flesh, and by the bones also of the vertebral tube. It is in fact easy to comprehend that the current of electric fluid, occasioned by the two coatings, penetrating only to a certain depth the parts of the animal covered by these coatings, can hardly reach the spinal marrow, or the principal branches of the nerves that enter into the interior parts of the limbs, if the bones, flesh, and other intervening integuments are of considerable

siderable thickness. The reason also must be obvious, why, in the larger animals, as dogs, lambs, &c. we fail to excite contractions in all the limbs by the application of the two coatings to the back, although stripped of its flesh. The large trunks of the nerves remain still at too great a depth; and it is only the smaller branches or ramifications that lay but a little below the coatings, and these branches terminate, for the most part, only in the neighbouring external parts; consequently we see produced only superficial contractions or palpitations in one or other of the muscles: or if by chance a whole limb is put in motion, it is because the nerve that goes to it, and influences this motion, is but thinly covered, so that only a thin layer of fibres intervenes between it and one or other of the metallic coatings, as appears from Experiment D. and the following ones (Sect. 23. &c.) in which the application of one of the coatings near the ischiatic nerve, in a dog or a lamb, was sufficient to excite considerable movements in the leg; and the nearer the coating was to the nerve, and the thinner the layer of flesh was that surrounded it, so much stronger in proportion were the contractions of the limb.

(44.) It becomes therefore necessary to know the situation of the nerves, their direction, &c. ; and it is requisite to remove not only the common integuments, the fat, &c. but likewise part of the flesh that covers and surrounds the nerves, in order that this surrounding muscular substance may be more or less extenuated, previously to the application of the metallic coating, to enable us to obtain in the larger animals contractions in any particular limb, to say nothing of the superficial contractions and palpitations of one or more muscles. It is perhaps impossible to excite these same motions and contractions in all the limbs at once ; although this is not difficult in the smaller animals, as we have already seen, (Sect. 42. Experiments O. and P.) merely by depriving them of the skin or a part of the other integuments ; and even this is not necessary in frogs, for in these animals we may leave the skin, it being so extremely thin and moist, as not to prevent, by its interposition, the electrical current from reaching the principal nerves or the spinal marrow.

(45.) But if it be necessary to pay attention to the direction of the principal nerves, in order to bring on the contractions in the different limbs, it is not less so to be careful of the position

sition of the coatings relatively to the muscles; for those muscles which are nearest to one or other of the coatings, are in general the most liable to contract spasmodic convulsions, and are oftentimes the only ones in which such an effect takes place; as, for instance, when the coatings do not correspond with any considerable nerve, or if there be a nerve, when it is surrounded with too much muscular flesh, or is too deeply seated.

(46.) This, and the Experiments E. F. (Sect. 28.) where a single muscle, and even a part of a muscle, treated in the usual way, experienced very strong contractions, might lead to a supposition that the electric fluid produces these effects by irritating the muscular fibres themselves, without the intervention of nerves; the action of which would consequently be neither *primary*, nor absolutely necessary, as I pretend. But an argument of this sort, founded on these facts, can have no weight, unless it could be proved that in these muscles, or portions of muscles, there are no nerves; for if there are nerves, (and certainly there must be, and are, nervous filaments in every sensible portion of a muscle, I had almost said in every muscular fibre) I may still maintain that it is these ner-

vous filaments, ramifying through the whole substance of a muscle, that are immediately affected by the electric fluid which penetrates this same substance ; that this fluid exerting its influence on their nerves, an influence that finishes there, the latter exert theirs on the muscles, &c. I may, I say, be able to maintain, with sufficient probability, that the electric fluid has no other influence, in the phenomenon of muscular contractions, than that of exciting the nerves ; in a word, that it is not the immediate cause. Such an assertion, which the things already explained render more than probable, is proved directly, and in the most obvious manner, by several experiments I have made on the tongue ; experiments that have led me to other discoveries equally interesting and curious.

(47.) Having succeeded in exciting tonic convulsions, and the most violent motions in the muscles and limbs, not only of small but of large animals, without laying bare any nerves, by the simple application of coatings of different metals to the muscles when freed from their integuments, I soon thought of trying whether the same effects might not be obtained in the human body. I conceived that the thing might succeed very well in amputated limbs ; but in the entire

tire

ture and living subject how was it to be effected? It seemed likewise to be necessary to remove the integuments, make deep incisions, and even dissect off portions of the flesh from the parts on which we might think of applying the metallic coatings (as I have remarked we are often obliged to do in the larger animals). Fortunately it came into my head, that we have, in the tongue, a muscle that is bare, or at least destitute of those thick integuments with which the external parts of the body are covered, a muscle which is extremely moveable, and moveable at will. Here then, I said to myself, are all the conditions requisite to enable us to excite movements by the usual artifice of different metallic coatings. With this view I made, on my own tongue, the following experiment.

(48.) EXPERIMENT Q. Having covered the point of the tongue, and a part of its upper surface, to the extent of some lines, with a piece of tin foil, (what is called silver paper is the fittest for the purpose) I applied the convex part of a silver spoon farther on, on the flat part of the tongue, and by inclining the spoon downwards brought the handle of it into contact with the tin foil. I expected to see my tongue affected with tremor; and on this account

count I made the experiment before a looking-glass. The effect, however, I had ventured to foretel did not take place; but instead of it I had a sensation I nowise expected; this was a pretty strong acid taste on the point of the tongue.

(49.) I was at first much surpris'd at this; but upon reflecting a little on the fact, I easily conceived, that the nerves which terminate on the point of the tongue; being the nerves destined for the sensations of taste, and not for the motion of this flexible muscle, It was perfectly natural, that the irritation of the electric fluid, put in motion by the usual artifice, should excite a taste, and nothing more; and that in order to excite in the tongue the motions of which it is susceptible, it would be necessary to apply one of the metallic coatings near its root, where the nerves enter that influence its motion; and this I soon verified by another experiment, as follows:

(50.) EXPERIMENT R. Having cut out, from a lamb recently killed, the tongue near its root, I applied a piece of tin foil at the end that was cut, and the silver spoon to one of the surfaces of the tongue; and then forming a communication between these two metallic coat-

ings, I had the pleasure to see the whole tongue affected with tremor, raising its point, and turning and bending itself in different directions, every time, and as long as such a communication took place.

(51.) I have repeated this experiment on the tongue of a calf, which I placed, coated in the same manner with a piece of tin foil near its root, on a silver plate, that the latter might serve as another coating; and the success was the same. I have likewise repeated it on the tongue of other smaller animals, as mice, chicken, rabbits, &c. and I have almost always obtained the same effect. I say *almost always*, for in the tongue of the smaller animals it sometimes failed; either because the tin foil was not applied exactly to the proper place, where the nerves that influence the motions of the tongue are inserted; or because the tongue being cold, had lost its vitality, which seldom lasts long in the muscles of animals of warm blood, as I have already had occasion to observe (Sect. 26.), and particularly in the tongue.

I am, &c.

A. VOLTA.

October 25, 1792.

XII. *A Return*

XII. Return* of the Ship's Company, and of the Military, on Board the Ships in the Service of the Honourable the United East-India Company, for the Years 1792 and 1793. By John Lorimer, M. D.

Ships' Names.	Destination.	Ships' Company outward.	Sick.	Re-covered.	Dead	Ships' Company homeward.	Sick.	Re-covered.	Dead	In Port.	Sick.	Re-covered.	Dead	Number of Recruits outward	Sick.	Re-covered.	Dead	Invalids, &c. homeward.	Sick.	Re-covered.	Dead	
Ocean	St. Helena, Madras, and China	134	102	100	1	124	45	45	0	St. Helena, Madras, and China	77	74	3	309	62	61	1	8	1	1	0	
Nottingham	Coast and China	128	67	65	2	112	12	12	0	Coast and China	54	44	10	218	100	106	3	52	0	0	0	
Lord Macartney	Ditto	105	49	49	0	105	22	21	1	Ditto	80	78	2	181	144	140	4	21	8	8	0	
Ganges	Ditto	106	40	33	7	104	14	14	0	Ditto	25	24	1	163	69	64	5	70	0	0	0	
Sir Edward Hughes	Ditto	No Regular Return out.				110	56	53	3	Ditto	No Return.				No Return.				5	0	0	0
Europa	Ditto	103	86	85	1	106	136	132	4	Ditto	148	126	5	428	264	263	1	65	45	42	3	
Melville Castle	Coast and Bay	106	15	13	1	103	25	22	3	Coast and Bay	38	30	5	337	125	119	1	39	12	10	3	
Contractor	Ditto	105	53	53	0	100	50	47	3	Ditto	89	82	7	193	186	183	3	82	53	48	5	
Ponshorne	Ditto	106	47	45	0	99	12	10	0	Ditto	59	58	1	157	22	20	2	98	5	4	1	
Busbridge	Ditto	106	56	55	1	94	12	11	1	Ditto	30	26	4	138	74	72	2	46	2	2	0	
Rose	Ditto	104	25	25	0	121	42	40	2	Ditto	39	37	2	142	35	32	3	88	13	11	2	
King George	Ditto	99	19	19	0	120	25	23	2	Ditto	26	24	2	155	26	24	2	69	13	11	2	
Rockingham	Bombay and China	106	42	41	1	101	9	8	1	Bombay and China	58	55	3	168	68	67	1	25	0	0	0	
Sullivan	Ditto	113	51	49	0	103	94	93	1	Ditto	89	82	7	95	39	39	0	10	3	3	0	
Middlesex	Ditto	111	20	15	5	107	10	9	1	Ditto	35	31	4	160	80	76	4	6	2	2	0	
Duke of Montrose	Bombay	101	46	46	0	101	31	31	0	Bombay	61	61	0	186	60	59	1	76	18	18	0	
General Elliot	Ditto	105	16	16	0	105	29	26	3	Ditto	47	43	4	176	26	24	2	61	3	1	2	
Earl of Wycombe	Bencoolen and China	87	83	82	1	82	27	24	3	Bencoolen and China	20	20	0	8	0	0	0	1	0	0	0	
Valentine	St. Helena, Bengal, and Bencoolen	106	19	19	0	97	30	27	3	St. Helena, Bengal, and Bencoolen	86	84	2	73	23	23	0	96	50	45	5	
General Goddard	Bengal	106	27	26	1	104	31	29	2	Bengal	71	61	10	155	110	109	1	64	14	14	0	
Bellmont	Ditto	90	110	110	0	85	65	60	5	Ditto	106	89	17	17	2	2	0	52	30	29	1	
Earl Talbot	Ditto	90	40	36	4	73	37	36	1	Ditto	27	25	2	15	2	1	1	28	10	7	3	
Lafcelles	China	108	67	67	0	108	11	8	3	China	116	110	6	4	0	0	0	7	0	0	0	
Walpole	Ditto	105	53	52	1	107	43	41	2	Ditto	73	70	3	0	0	0	0	2	0	0	0	
Thetis	Ditto	106	95	94	1	105	136	132	4	Ditto	75	73	2	0	0	0	0	6	0	0	0	
Royal Admiral	Port Jackson and China	121	25	24	1	120	54	33	3	Port Jackson and China	4	1	5	Convicts				13	3	0	0	
		2657	1253	1219	28	2701	1058	987	51		1533	1408	96	3929	1751	1623	50	1075	282	256	27	
Dublin	Of these three Ships, which failed in the Year 1791, there are no returns.																					
Airly Castle																						
Bridgewater																						

* In this Table the number of sick sometimes exceeds the number of the ship's company. If the same man is two or three times sick during the voyage, and recovers, this must be the case. Sometimes also the number of recovered and dead are not equal to the number of sick; but this must also happen when, at the end of the voyage, some still remain on the sick list.

XIII. *An Account of a singular Case of Ischuria, in a young Woman, which continued for more than three Years; during which Time, if her Urine was not drawn off with the Catheter, she frequently voided it by vomiting; and, for the last twenty Months, passed much Gravel by the Catheter, as well as by vomiting, when the Use of that Instrument was omitted, or unsuccessfully applied. To which are added some Remarks and Physiological Observations. By Isaac Senter, M. D. Associate Member of the College of Physicians of Philadelphia, and senior Surgeon in the late American Army. Vide Transactions of the College of Physicians, of Philadelphia. Vol. I. Part I. 8vo. Philadelphia, 1793.*

THE subject of this extraordinary case was a healthy-looking servant girl, who, in June, 1785, being then in her fifteenth year, was seized with a pain in the left hypochondrium, accompanied with cough, oppression at her breast, dyspnœa, and fever.

She had menstruated pretty regularly from the age of thirteen till within five weeks of her present illness, which was ascribed to cold.

Venaſection and other ſuitable remedies were had recourſe to by Dr. Senter, to whom ſhe applied for relief, and her complaints ſoon ſubſided; but about a month afterwards ſhe vomited up a quantity of bloody pus, which induced him to think a vomica had burſt in her ſtomach; for during the whole of this illneſs, her ſtomach, it ſeems, was ſo irritable, that ſhe could with difficulty retain in it either food or medicine.

She had now a ſuppreſſion of urine, which, after continuing twenty-four hours, went off without any medical aſſiſtance. After this ſhe became regular in her menſes, and in about two months was ſufficiently recovered to reſume her employment as a ſervant, which ſhe continued to follow till the 3d of June, 1786, when all her former complaints (except the ſuppreſſion of the menſes) returned with greater ſeverity than before.

Her pulſe was now at 120; her ſtomach, as during the former attack, was ſo irritable, that ſhe vomited up immediately almoſt every thing ſhe took. Of the different remedies that were had recourſe to, opium, when ſhe could retain it on her ſtomach, and repeated blood-letting in ſmall quantities, gave her the moſt relief.

On

On the 2d of July, when the severity of the symptoms had subsided, she was seized with a total suppression of urine, which continued till the beginning of the sixth day, when a vomiting came on, which lasted till she brought up nothing but water; and this water, she said, tasted like urine.

As the vomiting continued she found relief from the soreness and swelling she had felt for several days in the lower part of the abdomen.

She now thought herself much better, but the vomiting continued to return, more or less, every day, till the 14th of July, when Dr. Senter again saw her, and prevailed on her to submit to the introduction of a catheter, by means of which he drew off about three pints of clear, but high-coloured, urine.

From this time, till December, she continued with very little abatement of her complaints; and as she could lie in no other position, was constantly supported in an arm chair, in a reclined posture, with pillows under her hips.

During the whole of this period, whenever her water was omitted to be drawn off once in thirty or thirty-six hours at farthest, she never failed, we are assured, to vomit it up. To as-

certain so extraordinary a fact, our author tells us he often visited her about the time he knew she must vomit if the catheter was not introduced; and after examining her bladder, and finding it full, hard, and tender, sat by her till the vomiting returned, saved the water that she brought up in this way, and on comparing it with what he drew off by means of the catheter, found it the same in every respect.

During the time her urine came off by vomiting, she suffered, it seems, great anxiety and thirst, and complained of a sensation of inversion or turning up of something (running, as she expressed it) that appeared to tear her bowels.

In January, 1787, from some cause unknown, she could not be relieved with the instrument, nor could she vomit up her urine for several days; but at length it passed by the navel for three days successively; after which the catheter was used with the same effect as before.

About the beginning of August a brick-coloured gravel began to pass off through the catheter, and continued to be discharged in considerable quantity, whenever her urine was drawn off, till the beginning of November; at which time she felt more distress than usual, when-

whenever her urine came off by vomiting, and she soon observed a gritty substance in her mouth. When our author was informed of this new phenomenon, he requested her to save the urine for his inspection the next time she vomited; and on comparing it with what he drew off, found it contained the same kind of gravel as that which passed the catheter.

From this period, to the summer of 1788, her complaints, he observes, continued much the same; but during that summer she twice passed a small quantity of urine through the urethra, each time in consequence of being frightened. The hypogastrium became more tumid, and she complained of great soreness about the bladder, even after it was evacuated; the bladder itself seemed to be much thickened, and the apparent inequality of its surface was so great, and the tumour sometimes shifted so towards the right or left inguen, according as her body was moved, that our author suspected the existence of a stone.

Through the month of September her urine, we are told, could very rarely be drawn off; for upon the introduction of the catheter, a spasm seized the urethra and neck of the bladder, so that although the instrument seemed to

pass high up into the bladder, not more than a gill of urine could be drawn off, before it stopped entirely, with a sensation of something falling down against the cervix, which she was confident was a stone; and early in the following month, Dr. Senter being able to introduce a sound, readily met with a stone, which seemed to be of a small size, and softer than urinary calculi commonly are.

She had at different seasons of the year several small abscesses on different parts of her body, but they did not appear to relieve her general complaints. She also voided at times (after she had thrown up her urine) a bloody pus, of a coppery taste. This purulent discharge, it is observed, was never expectorated by coughing, though she had at times a dry cough, but was constantly brought up by vomiting.

In the spring of 1789 her urine began to pass *per anum*, loaded with the same kind of gravel that had come away by the catheter. This diminished but did not put a stop to her vomiting; for she continued to throw up more or less gravel that way every week. This new course of her urine occasioned a troublesome diarrhœa and tenesmus, but she felt less inconvenience from the stone in the bladder.

After

After the 13th of May her bladder never became so much distended with urine as it had been before; and the secretion of urine, as well as the formation of gravel, we are told, evidently diminished in proportion to her loss of strength, and the increase of the diarrhoea. The menses, which, during the whole of her illness, had returned at irregular periods, now entirely ceased. During the summer, the frequency of vomiting increased; she had several convulsive fits after vomiting; became more and more emaciated, and hectic; and, at last, lethargic; and on the 11th of August, 1789, died.

The body was examined the day after her death, by Dr. Senter, in the presence of Dr. Waterhouse, of Cambridge, and Dr. Mason, of Philadelphia, who, as well as several other respectable medical practitioners, had occasionally visited her in her life-time, and seen her vomit up both urine and gravel.

On dissection, nothing was discovered that could throw any light on the nature of the disease.

In the thorax, the only morbid appearance was an adhesion of part of the right lobe of the lungs to the pleura.

In

In the abdomen, the omentum was found much wasted, and of a dark gangrenous colour; the stomach also is described as being in a gangrenous state, and containing ‘a semi-purulent matter, of a fœtid scent;’ but the weather, we find, was very warm, and the body in an offensive state, at the time the dissection was made. Nothing particularly worthy of notice was observed in the state of the liver, gall-bladder, intestines, kidneys, or ureters. The urinary bladder was also in its natural state, not in the least thickened, and contained no sand or gravel. The uterus contained about a drachm of thick, fœtid pus, but had no other appearance of disease; the Fallopian tubes were larger than usual, and strung with several hydatids of the size of a walnut; the corpora fimbriata had a gangrenous appearance; the ovaria were enlarged to the size of a small hen’s egg, and distended with a clear limpid fluid.

To the preceding history Dr. Senter has added many judicious remarks; and in his attempt to account for the phenomena of so very uncommon a case, has not omitted to avail himself of the modern doctrine of the retrograde motion of the lymphatics, and of the opinions of those writers who have maintained
the

the existence of a direct communication between the alimentary canal and the urinary bladder.

There are many instances, he observes, in medical books, of sudden and partially-increased actions of the vessels of the human body; but he candidly acknowledges that his reading has furnished him with no fact similar to the extraordinary one which is the subject of the paper before us*: that which he considers
as

* There are, however, upon record, two cases which exhibit a striking analogy to that of Dr. Senter's patient; and although they may have been overlooked, or perhaps disregarded on a supposition of their improbability, they must now become extremely interesting by the tendency they have to corroborate the curious and extraordinary facts he has related. Both the cases we allude to occur in the History of the Academy of Sciences at Paris, and are as follows:

Case I. " M. Maraldi has communicated to the Academy the following case, from a letter addressed to him by M. Marangoni, physician at Mantua:

" A Nun, of the Order of St. Francis, in the convent of St. Joseph, at Mantua. aged thirty-five years, of a thin and delicate habit of body, and who had long been subject to hysterical complaints, was attacked with pains, spasms, and swelling of the abdomen, to which succeeded a violent and alarming suppression of urine. Soon after this she felt a pain, which she described as ascending from the lower part of the abdomen to her stomach; and she
" vomited

as coming the nearest to it, is a case described by Dr. Percival, in the second volume of his *Essays, Medical and Experimental*, (8vo, London,

“ vomited a fluid which, without any difficulty, was
 “ known to be urine. This vomiting continued forty days,
 “ during which time the patient voided no urine by the
 “ usual channel, unless the surgeon drew it off with a ca-
 “ theter, and even then the quantity scarcely amounted to
 “ an ounce a day. At the end of the forty days, the urine
 “ spontaneously resumed its natural course, and in a day or
 “ two the patient found herself perfectly recovered. But
 “ the vomiting of urine returned, and at the end of twenty-
 “ seven days, the patient complained of very acute pain
 “ about the region of the pubis. Her surgeon was desirous
 “ of relieving her by means of the catheter, but there was
 “ such a contraction of the urethra, that he found it impos-
 “ sible to introduce even a probe into the bladder. The
 “ vomiting of urine has continued, and what is remarka-
 “ ble, there is no appearance of food mixed with it, even
 “ when the vomiting takes place soon after her meals.
 “ When M. Marangoni wrote this account, the patient had
 “ been in this state thirty-two days.

“ This singular complaint would lead one to think there
 “ is an immediate though hitherto undiscovered commu-
 “ nication between the stomach and the urinary bladder;
 “ but M. Marangoni and the celebrated Lancisi are of a
 “ different opinion; they both of them think, that in cases
 “ of this kind a suppression of urine takes place in the kid-
 “ neys; that is to say, that the kidneys cease to extract this
 “ fluid

London, 1773) of a woman who, after a spontaneous vomiting of several days, during which she brought up three gallons of water, was entirely cured of a dropfy of the ovarium.

“ fluid from the blood, and that in their stead the glands of
“ the stomach perform this function.”

Case II. “ M. Lemery is acquainted with a Monk, who,
“ for about eight years, has been subject to a periodical vo-
“ miting, the fits of which are as regular as those of a quar-
“ tan ague. Five hours, or thereabouts, before the vomit-
“ ing begins he complains of violent pains in his kidneys.
“ The vomiting continues, with intervals, four or five
“ hours. What he vomits is of a dirty red colour. It is
“ almost entirely water, but has a strong urinous smell, and
“ the patient has no doubt of its being really urine, as he
“ eats but very little, and drinks more than the usual portion
“ of a Monk. He drinks only wine, the colour of which
“ agrees with that of the fluid he vomits. A few hours
“ after the vomiting he finds himself well, and remains so
“ till the next fit. He uses a great deal of exercise, with-
“ out which he thinks he should suffer more. It is a known
“ fact, that in nephritic pains, which are always occasioned
“ by obstructions of the kidneys, the patients are subject to
“ frequent vomiting, and that what they bring up smells
“ much of urine.”—*See Histoire de l'Academie Royale*
des Sciences, Années 1715 & 1722. EDITOR.

CATALOGUE OF BOOKS.

1. **T**HOUGHTS on the Effects of the Application and Abstraction of Stimuli on the Human Body ; with a particular View to explain the Nature and Cure of Typhus. By *J. Wood*, M. D. 8vo. *Murray*, London, 1793.

2. An Account of the Bilious, Remitting, Yellow Fever, as it appeared in the City of Philadelphia in the year 1793. By *Benjamin Rush*, M. D. 8vo. Philadelphia, 1794.

3. Observations on the Cause, Nature, and Treatment of the Epidemic Disorder prevalent in Philadelphia. By *D. Naffy*, M. D. Member of the American Philosophical Society. 8vo. Philadelphia, 1793.

4. A Short Account of the Malignant Fever, lately prevalent in Philadelphia ; with a Statement of the Proceedings that took place on the Subject in different Parts of the United States. By *Matthew Carey*. 8vo. Philadelphia, 1793.

5. A Treatise on the Extraction of the Cataract. By *Frederick Bischoff*, F. M. S. Oculist to his Majesty in the Electorate of Hanover,

and to her Majesty in England. 8vo. *Nicol*, London, 1793.

6. An Account of a Fever which appeared in several Parts of Somersetshire in the year 1792. By *Richard Poole*, Surgeon, Sherborne. 8vo. *Johnson*, London, 1793.

7. A Guide for Self-Preservation and Parental Affection; or plain Directions for enabling People to keep themselves and their Children free from several common Disorders. By *Thomas Beddoes*, M. D. 12mo. *Murray*, London, 1793.

8. A Chemical Dissertation on the Thermal Waters of Pifa, and on the neighbouring acidulous Spring of Asciano; with an Historical Sketch of Pifa, and a Meteorological Account of its Weather. To which are added, Analytical Papers respecting the Sulphureous Water of Yverdun. By *John Nott*, M. D. of Bristol Hot-wells. 8vo. *Walter*, London, 1793.

9. Horti Botanici Cantabrigienfis Catalogus. 8vo. Cantabrigiæ, 1794.

10. Flora Oxoniensis, exhibens Plantas in agro Oxoniensi sponte crescentes, secundum Systema sexuale distributas. Auctore *Joanne Sibthorp*, M. D. Professore Regio Botanico, Regiæ Societatis Londinensis aliarumque Societatum Socio. 8vo. Oxonii, 1794.

11. Differ-

11. Differtatio Inauguralis de Angina maligna. Auctore *Arthuro Bedford*, Anglo. 8vo. Edinburgi, 1792.

12. Differtatio Inauguralis de Respiratione. Auctore *Thoma Blair*, Scoto-Britanno. 8vo. Edin. 1792.

13. Differtatio Inauguralis de Variolis. Auctore *Joanne Bower*, Scoto. 8vo. Edin. 1792.

14. Differtatio Inauguralis de Visu. Auctore *Wheaton Bradish*, Hiberno. 8vo. Edin. 1792.

15. Differtatio Inauguralis de Rheumatismo acuto. Auctore *Joanne Bradley*, Anglo. 8vo. Edin. 1792.

16. Differtatio Inauguralis de Cœli Effectibus. Auctore *Jacobo Buchan*, Scoto. 8vo. Edin. 1792.

17. Differtatio Inauguralis de Rheumatismo acuto. Auctore *Andrea Grieve*, Scoto. 8vo. Edin. 1792.

18. Differtatio Inauguralis de Hypochondriaci. Auctore *David Corbin Kerr*, Virginienſe. 8vo. Edin. 1792.

19. Differtatio Inauguralis de Variolis. Auctore *Gulielmo Marsden*, Anglo-Britanno. 8vo. Edin. 1792.

20. Differtatio Inauguralis de Pneumonia.

Auctore *Carolo Merivether*, Virginienſe. 8vo.
Edin. 1792.

21. Differtatio Inauguralis de Variolis. Auc-
tore *Roberto Montgomery*, Hiberno. 8vo. Edin.
1792.

22. Differtatio Inauguralis de Hydrope Ana-
ſarca. Auctore *Thoma Pollard Pierce*, Barba-
denſe. 8vo. Edin. 1792.

23. Differtatio Inauguralis de Angina ma-
ligna. Auctore *Georgio Wier*, Scoto. 8vo
Edin. 1792.

24. Differtatio Inauguralis de Alimento.
Auctore *Gulielmo Yates*, Anglo. 8vo. Edin.
1792.

25. Differtatio Inauguralis de Coitu ejusque
variis Formis quatenus Medicorum ſunt. Auc-
tore *Johanne Paul Gottlieb Kircheiſen*. 4to.
Jena, 1792.

26. Analyſe du Syſteme abſorbant ou lym-
phatique. Par *M. des Genettes*, D. M. 8vo.
Montpellier, 1791.

27. Memoire ſur une Maladie de l'Ovaire.
Par *Jean Baptiſte Ph. R. N. Laumonier*, Chirur-
gien en chef de l'Hotel Dieu de Rouen. 4to.
Rouen, 1790.

28. Avis aux Sages Femmes; par *M. Sacombe*,
Vol. VI. Q Medecin

Medecin-Accoucheur, Membre de plusieurs Academies. 8vo. Paris, 1792.

29. Recherches Physico-chymiques. Cahiers I. II. III. 4to. Amsterdam, 1793-4.

30. Memoires de l'Academie Royale des Sciences et Belles Lettres depuis l'Avenement de Frederic Guillaume II. au Trone. 1788 et 1789. Avec l'Histoire pour le même Tems. 4to. Berlin, 1793.

31. Sammlung der Deutschen Abhandlungen welche in der Königlichen Akademie der Wissenschaften zu Berlin vorgelesen worden in den Jahren 1788 und 1789. *i. e.* A Collection of German Essays, read before the Royal Academy of Sciences at Berlin, in the Years 1788 and 1789. 4to. Berlin, 1793.

32. Memoria Chirurgica sul Labbro leporino complicato; di *Giuseppe Sonfis*, R. Affess. Medico, &c. 4to. Cremona, 1793.

33. Pisaura Automorpha e Coreopsis formosa; Piante nuove pubblicate da *Giuseppe Antonio Bonato*, Dott. di Medicina, pubblico Bibliotecario, Ispettore e Soprantendente all' Orto medico dell' Università di Padova. 4to. Padova, 1793.

I N D E X.

A.

A CID, Vegetable, Solution of Sal Ammoniac in, good effects of, in lacerated Wounds, —	66
Africa, Coast of, Observations on the Diseases most frequent there, — — —	60
Aliment, Dissertation on, ———	226
Ammoniacal Salt, Solution of, in Vinegar, good effects of, in lacerated Wounds, ——— ———	66
Anasarca, Dissertation on, ———	226
Aneurism, of the Crural Artery, Case of, —	114
Angina, malignant, Dissertations on, —	225, 226
Animal Electricity, Observations and Experiments relative to, ——— ———	161
Antimony, tartarised, recommended as the best Emetic in the Intermittents of Tropical Climates, ———	42
———, ill effects of nauseating Doses of, during the State of apyrexia in such Cases, ———	<i>ibid.</i>
Arsenic, Observations on its Use in Intermittents, 1, 46, 61	
———, Effects of, in such Cases, when combined with Peruvian Bark, ——— ———	43
———, said to be as powerful and nearly as certain as the Bark, in the Cure of Intermittents; but allowed to be inferior to the Bark in its tonic Effects, —	46
———, Conjecture concerning its specific Action, and that of the Bark, in the Cure of Intermittents, ———	47
———, Account of irregular Cases of Intermittents in which it fails, ——— ———	44
———, Historical Account of its Use in Intermittents, ———	46
Atmospheres, Electrical, Effects of, on Frogs and other Animals, ——— ———	166

B.

Bark, Peruvian, the only Remedy to be relied on in the Intermittents of Tropical Climates threatening immediate Danger, ——— ———	44
———, Conjectures relative to its specific Action in the Cure of Intermittents, ———	47, 143
Q 2	Bark,

Bark, Peruvian, Prejudices which still prevail against its Use,	57
———, different Species of, enumerated,	146
Beddoes, Thomas, Guide for Self-preservation,	224
Bedford, Arthur, de Angina maligna,	225
Berlin, Memoirs of the Academy of Sciences at,	227
Bischoff, Fred. on the Extraction of the Cataract,	223
Blair, Thomas, de Respiratione,	225
———, William, of the Extraction of an extraneous Substance from the Rectum,	111
Bonato, G. Anton. Description of two new Plants,	227
Bower, Joannes, de Variolis,	225
Bradish, Wheaton, de Vifu,	<i>ibid.</i>
Bradley, Joannes, de Rheumatismo acuto,	<i>ibid.</i>
Bread, toasted, Instance of its producing painful Effects in the Rectum,	111
Buchan, Jacobus, de Coeli Effectibus,	225

C.

Cambridge, Catalogue of the Botanic Garden at,	224
Camper, Professor, his <i>Demonstr. Anat. Pathol.</i> referred to,	107, 108
Carey, Matthew, Account of the Yellow Fever at Philadelphia,	223
Carter, Henry Yates, on the Effects of Sal Ammoniac and Vinegar on lacerated Wounds,	66
———, Case of a diseased Kidney,	85
———, Gun-shot Wound,	91
Cataract, Treatise on the Extraction of,	223
Caterpillars, not affected by Electricity,	193
Cinchona, different Species of, enumerated,	146
Clarke, Robert, Account of a new Key Instrument,	120
Climate, Dissertation on the Effects of,	225
Coating, Metallic, Advantages of, in Experiments of Animal Electricity,	174
———, of two different Metals, Observations on the Use of, in Experiments of Animal Electricity,	178, 183

D.

Diarrhœa, Effects of Mahogany Wood in Cases of,	156
Duncan, Dr. Andrew, Junior, his Dissertation on the Swietenia Soyimida referred to,	127, 144

E.

Electricity, Animal, Observations and Experiments relative to,	161
Electricity,	

Electricity, Discovery of a new Law in, —	177, 192
———, Estimate of the Quantity of, necessary to produce certain Effects in a Frog, —	170
Electrometer, Animal, what so called, —	171
Emetics, good Effects of, in Intermittents, —	41, 42

F.

Fever, Epidemic, in Somersetshire, Work relative to, —	224
———, Intermittent, of a Tropical Climate, Observations on, —	1
———, ————, how distinguished from the Remittent, —	43
———, ————, anomalous Cases of, described, —	44
———, ————, See <i>Antimony</i> , <i>Arsenic</i> , <i>Bark</i> , <i>Emetics</i> , <i>Opium</i> .	

———, Intermittent and Remittent, the most frequent Diseases on the Coast of Africa, —	60
———, Yellow, at Philadelphia, Works relative to, —	223
Forster, Thompson, Case of Aneurism of the Crural Artery, —	114
Frogs, Effects of Electric Atmospheres on, —	166

G.

Galvani, Lewis, Account of his Discoveries relative to Animal Electricity, —	161
Genettes, M. des, Analyse du Systeme Lymphatique, —	226
Gravel, voided, with Urine, by vomiting, —	215
Grieve, Andreas, de Rheumatismo acuto, —	225

H.

Hare Lip, Work relative to, —	227
Head, Case of a Gun-shot Wound of, —	91
Hughes, Francis, on the Effects of Mahogany Wood in Diarrhoea, —	156
Hypochondriacal Affection, Dissertation on, —	225

I.

Insects, several Species of, not affected by Electricity, —	194
Intermittents of a Tropical Climate, Observations on the Use of Arsenic in, —	1
———, ————, Use of Opium in, —	41
———, ————, Emetics in, <i>ibid.</i>	
———, ————, how distinguished from a Remittent Fever, —	43
———, ————, Circumstances of, in which the Bark alone is to be relied on, —	44
———, ————, Intermittents	

Intermittents of a Tropical Climate, anomalous Cases of, described, _____	_____	<i>ibid.</i>
Ischuria, singular Case of, _____	_____	212
Issue, singular Effects of, _____	_____	109

K.

Kerr, David Corbin, de Hypochondriasi, _____	_____	225
Key Instrument, of a new Construction, Account of, _____	_____	120
_____, Observations on the Principles on which it acts, _____	_____	124
Kidney diseased, Case of, _____	_____	85
_____, Appearance of, on Dissection, _____	_____	89
Kircheisen, I. P. G. Dissertatio de Coitu, _____	_____	226

L

Lancisi, his Theory of a Case in which Urine was voided by vomiting, _____	_____	221
Laumonier, M. sur une Maladie de l'Ovaire, _____	_____	226
Lemery, M. Case of a Monk who vomited Urine, _____	_____	222
Leyden Phial, the supposed Analogy of, to some Phenomena of Animal Electricity, _____	_____	190
Lizard, Experiment on, _____	_____	202
Lorimer, Dr. John, Return of the Sick on Board the East- India Company's Ships for the Years 1792 and 1793, _____	_____	211
Lymphatic System, Work relative to, _____	_____	226

M.

Mahogany, a new Species of, described, and its Bark recom- mended as a Substitute for the Peruvian Bark, _____	_____	127
_____, Wood, Account of its Effects in Diarrhœa, _____	_____	156
_____, of Jamaica, described as more astringent than that of Honduras, _____	_____	157
Marangoni, M. Case of a Patient who vomited Urine, _____	_____	220
Marfden, Gulielmus, de Variolis, _____	_____	225
Melia Azadirachta, the Bark of, recommended as a Substitute for Peruvian Bark, _____	_____	154
Merrivether, Carolus, de Pneumonia, _____	_____	225
Montgomery, Robertus, de Variolis, _____	_____	226
Muscles, Experiments on, _____	188, 191, 198,	202
_____, not <i>immediately</i> affected by Electricity, _____	_____	179, 200
_____, voluntary, the only ones affected by weak Currents of Electricity, _____	_____	197

N.

Nassy, Dr. D. Observations on the Epidemic Fever at Phila- delphia, _____	_____	223
		Naucleæ

Nauclea Daduga, Bark of, its Properties described,	155
Nerves, the only Parts immediately affected by Electricity,	179, 200
———, Effects of certain morbid Alterations of,	96
———, Experiments on,	181
———, painful Effects of Pressure on,	101
———, subcutaneous, a Disease of, described,	108
Nott, Dr. John, on the Waters of Pisa and Yverdun,	224

O.

Opium, its Effects in the Intermittents of a tropical Climate,	41
Ovarium, Dropsy of, cured by a spontaneous vomiting,	221

P.

Pearson, John, Account of extraordinary Symptoms arising from certain morbid Affections of the Veins and Nerves,	96
Percival, Dr. his Account of a Dropsy of the Ovarium cured by a spontaneous vomiting, referred to,	221
Philadelphia, Works relative to the Yellow Fever at,	223
Pierce, Thomas Pollard, de Hydrope Anasarca,	226
Pisa, Dissertation on the Thermal Waters of,	224
Pneumonia, Dissertation on,	225
Poole, Richard, Account of an Epidemic Fever in Somersetshire,	224
Pouteau, M. his Posthumous Works quoted,	103

Q.

Quadrupeds, Experiments on,	181, 187
-----------------------------	----------

R.

Recherches Physico-chymiques,	227
Remittent Fever, <i>See</i> Fever.	
Respiration, Dissertation on,	225
Rheumatism, acute, Dissertation on,	225
Roxburgh, Dr. W. Account of a new Species of Swietenia,	127
Rush, Dr. Benj. of the Bilious Remitting Yellow Fever,	223

S.

Sacombe, M. Avis aux Sages Femmes	226
Sal Ammoniac.— <i>See</i> Ammoniacal Salt.	
Senter, Dr. Isaac, of a singular Case of Ischuria	212
Sibthorp, Joannes, Flora Oxoniensis	224
Sierra Leone, Account of the Weather at,	2
Small Pox, Dissertations on,	225, 226
Sonfis, G. his Work relative to the Hare Lip	227
Stimuli, Work relative to the Effects of,	223
Swietenia,	

Swietenia, Accounts of two new Species of, 127, 153

T.

Teeth, Observations on the Extraction of, by the Key Instrument, 121

Tongue, human, Experiments on, 208

——, of Quadrupeds, Experiments on, 209, 210

Typhus, Work relative to the Nature and Cure of, 223

V.

Vinegar, Solution of Sal Ammoniac in, Effects of, on lacerated Wounds, 66

Vision, Dissertation on, 225

Volta, Alexander, Translation of his Letters on Animal Electricity, 161

Urine, Cases in which it has been voided by vomiting, 212, 220, 222

W.

Wallurfe, a Tree so called by the Hindoos, described, 154

Wier, Georgius, de Angina maligna, 226

Winterbottom, Dr. T. M. on the Use of Arsenic in Intermittents, 1

——, Account of the Weather at Sierra Leone, 2

Worms, not affected by Electricity, 194

Wood, Dr. J. on the Application and Abstraction of Stimuli, 223

Y.

Yates, Gulielmus, de Alimento, 226

Yellow Fever, Works relative to it, 223

Yverdun, Work relative to the sulphureous Water of, 224

